

# MAXIMUM PRINCIPLES FOR VECTORIAL MINIMIZERS OF NONCONVEX FUNCTIONALS IN $L^p$ AND $L^\infty$

Nikolaos I. Katzourakis\*

## Abstract

We establish *Maximum Principles* (MPs) which apply to *minimizers* and *absolute minimizers*  $U : \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^N$  of *integral* and *supremal functionals*  $E_\alpha(U, \Omega) = \|\mathcal{L}(\nabla U, U, \cdot)\|_{L^\alpha(\Omega)}$  for  $\alpha = 1$  and  $\infty$  respectively,  $n, N \geq 1$ . Our main result is  $U(\Omega) \subseteq \overline{\text{co}}(U(\partial\Omega))$ , suitably interpreted within our setting. We derive it by introducing a *new nonlinear convergence technique independent of weak\* LSC of  $E_\alpha$* . It is based on a  $O(N, \mathbb{R})$  invariance of  $\mathcal{L}$ , bypassing quasiconvexity. MPs are implied for minimal solutions to the non-elliptic *Euler & Aronsson* PDEs, previously known only in certain elliptic cases. If  $N = 1$ , we recover  $\sup_\Omega [\pm u] \leq \max_{\partial\Omega} [\pm u]$ . Our assumptions include the  $A$ - &  $\infty$ -Laplace systems  $\text{Div}(A(|\nabla U|)\nabla U) = 0$ ,  $(\nabla U)^\top [\nabla^2 U \cdot \nabla U] = 0$  and many stored energies of Elasticity. If existence fails for  $E_1$ , the MPs apply to minimizing sequences and to the relaxed Young measure solution. This work is an outgrowth of ODE ideas, first appearing in recent work with Alikakos [K-A].

## 1 Introduction.

The Maximum Principle (MP) is a major tool for the 2nd order elliptic PDEs. Roughly, the idea is that if  $u : \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$  solves  $F(x, u, \nabla u, \nabla^2 u) = 0$ , then  $\sup_\Omega [\pm u] \leq \max_{\partial\Omega} [\pm u] + C$  for  $C \geq 0$ . The MP has been initiated by Hopf [H]. For extensions, we refer to Puzzi-Serrin [P-S], Gilbarg-Trudinger [G-T], Protter-Weinberger [P-W], Fraenkel [Fr], Crandall-Ishii-Lions [C-I-L] and Jensen [Je1].

In the present paper we establish MPs for general vector functions  $U : \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^N \equiv \mathbb{R}^{1 \times N}$ ,  $U = (U_1, \dots, U_N)$ , which satisfy a minimality property with respect to the functional

$$E_\alpha(U, \Omega) := \|\mathcal{L}(\nabla U, U, \cdot)\|_{L^\alpha(\Omega)}, \quad \text{when } \alpha = 1 \text{ or } \infty. \quad (E_\alpha)$$

Here  $\Omega$  is open,  $\mathcal{L} : \mathbb{R}^{n \times N} \times \mathbb{R}^N \times \Omega \rightarrow \mathbb{R}$  is a Carathéodory function and the derivatives are understood as  $\nabla U = [\nabla U_1 | \dots | \nabla U_N] : \Omega \rightarrow \mathbb{R}^{n \times N}$ ,  $\nabla^2 U = [\nabla^2 U_1 | \dots | \nabla^2 U_N] : \Omega \rightarrow \mathbb{R}^{n \times nN}$ , etc.  $(E_\alpha)$  is associated to a 2nd order *PDE system* and our MPs apply to the PDE as well. Our approach is based only the minimality and not the solution property. Hence, we can derive MPs for  $(E_\alpha)$  while the PDE may have no solutions. We have written  $(E_\alpha)$  as a compact notation which includes two distinct kind of functionals. When  $\alpha = 1$ ,  $(E_\alpha)$  is the standard integral functional

$$E_1(U, \Omega) = \int_\Omega \mathcal{L}(\nabla U, U, x) dx \quad (E_1)$$

\*e-mail: nkatzourakis@math.uoa.gr, Department of Mathematics, University of Athens, Greece.

and the minimality is merely the minimizing property. Minimizers of  $(E_1)$ , if any, solve weakly (in a specified space setup) the system of *Euler* PDEs

$$\operatorname{Div}(\mathcal{L}_P(\nabla U, U, x)) = \mathcal{L}_\xi(\nabla U, U, x). \quad (\text{E})$$

Here the arguments of  $\mathcal{L}$  are denoted by  $\mathcal{L}(P, \xi, x)$ . If  $P = [P_{ia}] \in \mathbb{R}^{n \times N}$ , we set  $\mathcal{L}_P := ([\mathcal{L}_{P_{ia}}])^\top = [\mathcal{L}_{P_{ai}}]$ ,  $\mathcal{L}_\xi := \nabla_\xi \mathcal{L}$  and for  $A : \Omega \rightarrow \mathbb{R}^{N \times n}$  we set  $\operatorname{Div}(A)(x) := [\sum_i (A_{ai})_{x_i}]$ , with  $\operatorname{Div}(A) : \Omega \rightarrow \mathbb{R}^{N \times 1}$ . For direct methods for  $(E_1)$ , we refer to Dacorogna [D]. When  $\alpha = \infty$ ,  $(E_\alpha)$  is the supremal functional

$$E_\infty(U, \Omega) = \operatorname{ess\,sup}_{x \in \Omega} \mathcal{L}(\nabla U, U, x). \quad (E_\infty)$$

The functional  $(E_\infty)$  can be studied in  $[W^{1,\infty}(\Omega)]^N$  and  $[L^\infty(\Omega)]^N$ . When  $\alpha = \infty$  the correct minimality notion differs because in general minimizers of  $(E_\infty)$  do *not* solve an associated ‘‘Euler’’ PDE. The problem is that  $(E_\infty)$  is not local: minimizers of  $(E_\infty)$  on  $\Omega$  do not always minimize  $E_\infty(U, \Omega')$  on  $\Omega' \subset \subset \Omega$ . This does not arise in  $(E_1)$ . We rectify it by building locality in the definition of minimizers:  $U^* : \Omega \rightarrow \mathbb{R}^N$  is an *Absolute Minimizer* of  $(E_\infty)$  (usually written  $U^* \in \operatorname{AM}(E_\infty, \Omega)$ ) if

$$E_\infty(U^*, \Omega') \leq E_\infty(U^* + \phi, \Omega') \quad \text{for all } \Omega' \subset \subset \Omega \text{ and all } \phi \in [W_0^{1,\infty}(\Omega')]^N. \quad (\text{AM})$$

The variational property (AM) is the key for  $(E_\infty)$ . Absolute minimizers solve the *Aronsson* PDE system

$$\mathcal{L}_P(\nabla U, U, x) \nabla (\mathcal{L}(\nabla U, U, x)) = 0. \quad (\text{A})$$

In general, the highly singular PDE (A) has no  $C^2$  solutions ([Ar6]). When  $N = 1$ , AM’s of  $(E_\infty)$  are interpreted as viscosity solutions to (A). In certain cases, viscosity solutions to (A) verify (AM). We refer to Barron-Jensen-Wang [BJW1], Yu [Y] and to Crandall [Cr2] and Crandall-Wang-Yu [C-W-Y].

All MPs which apply to scalar  $u : \Omega \rightarrow \mathbb{R}$  invoke the ordering  $\leq, \geq$  of  $\mathbb{R}$ , but no unique analog exists when  $N > 1$ . We choose to substitute  $\leq, \geq$  by  $\subseteq, \supseteq$  with respect to convex sets  $C \subseteq \mathbb{R}^N$ . We derive pointwise MPs which do not involve norm bounds, being a direct extension of

$$\sup_{\Omega} u \leq \max_{\partial\Omega} u, \quad \inf_{\Omega} u \geq \min_{\partial\Omega} u. \quad (\text{MP-s})$$

Our viewpoint is based on the observation that (MP-s) can be recast as  $u(\Omega) \subseteq [\min_{\partial\Omega} u, \max_{\partial\Omega} u]$ . When  $N > 1$ , the appropriate vectorial extension of (MP-s) (equivalent to it when  $N = 1$ ) is

$$U(\Omega) \subseteq \overline{\operatorname{co}}(U(\partial\Omega)), \quad (\text{MP-v})$$

suitably interpreted (definitions (Co), (CO), (EssCo)) to make sense for merely measurable  $U$  and any  $\Omega$ . Hence, *the range is contained into the closed convex hull of the boundary values*.

The formula (MP-v) has appeared in the literature before for certain convex densities and elliptic systems. In Differential Geometry, a version of (MP-v) which reads  $\mathcal{M} \subseteq \overline{\operatorname{co}}(\partial\mathcal{M})$  is the well known ‘‘Convex Hull Property’’ of minimal submanifolds  $\mathcal{M} \subseteq \mathbb{R}^N$ . We refer to Osserman [Os1], [Os2], Colding-Minicozzi II [C-M1], [C-M2] and Alexander-Ghomi [A-G]. Recently, Bildhauer-Fuchs [B-F2] established (MP-v) for  $(E_1)$  in the broad  $[BV(\Omega)]^N$  setting of generalized minimizers, but for special  $\mathcal{L}$

of the form  $\mathcal{L} = \mathcal{L}(|P|)$  with  $p \mapsto \mathcal{L}(p)$  convex and increasing with  $\Omega$  a bounded Lipschitz domain. In the  $[W^{1,p}(\Omega)]^N$  setting, derivation of (MP-v) is trivial when  $\mathcal{L} = \mathcal{L}(|P|, x)$  with  $p \mapsto \mathcal{L}(p, \cdot)$  convex and increasing. Related material appears also in D'Ottavio-Leonetti-Musciano [D-L-M], Bildhauer-Fuchs [B-F1], Campanato [Ca], Cellina [Ce], Felmer-Quass [F-Q], Leonetti-Siepe [L-S], Leonardi [Le], Nečas-Stará [N-S], Alikakos-Fusco [A-F], [A-F2] and Katzourakis-Alikakos [K-A]. However, in all of these papers are considered situations either in a different context or in less generality. The present work extends among others the special MP tool developed in [K-A] for ODEs.

In this paper we develop a new nonlinear convergence technique which establishes (MP-v) without invoking the weak\* LSC of  $(E_\alpha)$ , hence we bypass quasiconvexity of  $\mathcal{L}$ . In both bases of  $(E_1)$  and  $(E_\infty)$ , such a notion is crucial for existence, which however conflict to each other. Let  $Q \equiv ((0, 1))^n$ . For  $(E_1)$ ,  $L : \mathbb{R}^{n \times N} \times \mathbb{R}^N \times \Omega \rightarrow \mathbb{R}$  can be called (Morrey)  $M$ -quasiconvex if for all  $(\bar{P}, \bar{\xi}, \bar{x}) \in \mathbb{R}^{n \times N} \times \mathbb{R}^N \times \Omega$ ,

$$L(\bar{P}, \bar{\xi}, \bar{x}) = \min \left\{ \int_Q L(\bar{P} + \nabla \phi(x), \bar{\xi}, \bar{x}) dx, \text{ over all } \phi \in [C_c^\infty(Q)]^N \right\}. \quad (\text{M-QC})$$

For  $(E_\infty)$ , we recall the definition of Barron-Jensen-Wang [BJW2]:  $L : \mathbb{R}^{n \times N} \times \mathbb{R}^N \times \Omega \rightarrow \mathbb{R}$  can be called for distinction  $BJW$ -quasiconvex, if for all  $(\bar{P}, \bar{\xi}, \bar{x}) \in \mathbb{R}^{n \times N} \times \mathbb{R}^N \times \Omega$ ,  $\varepsilon > 0$  and  $M > 0$ , there is a  $\delta = \delta(\varepsilon, M, \bar{P}, \bar{\xi}, \bar{x}) > 0$ : if  $\psi \in [W^{1,\infty}(Q)]^N$  satisfies  $\|\nabla \psi\|_{[L^\infty(Q)]^{n \times N}} \leq M$  and  $\|\psi\|_{[L^\infty(\partial Q, \mathcal{H}^{n-1})]^N} \leq \delta$ , then

$$L(\bar{P}, \bar{\xi}, \bar{x}) \leq \text{ess sup}_{x \in Q} L(\bar{P} + \nabla \psi(x), \bar{\xi}, \bar{x}) + \varepsilon. \quad (\text{BJW-QC})$$

Our method for  $(E_\alpha)$  bypasses (M-QC), (BJW-QC). It requires only on the invariance of  $P \mapsto \mathcal{L}(P, \cdot, \cdot)$  to reflections with respect to a convex set (assumption (HR)). In particular, it is automatically satisfied if  $\mathcal{L} = \mathcal{L}(|\nabla U|, U, x)$ . As a result, we derive MPs which apply to non-elliptic PDE systems (E) and (A). In the case of non-existence of minimizers for  $(E_1)$ , we derive (MP-v) for minimizing sequences and for the relaxed solution (equivalently, the Young measure solution) of the weak LSC envelope of  $(E_1)$ . Our assumptions include the  $\infty$ -Laplacian  $(\Delta_\infty)$  and all A-Laplacians (terminology of Puzzi-Serrin [P-S]):

$$\Delta_\infty U := (\nabla U)^\top [\nabla^2 U \cdot \nabla U] = 0, \quad \mathbf{A}U := \text{Div}(\mathbf{A}(|\nabla U|)\nabla U) = 0. \quad (\Delta_\infty\text{-A})$$

Here  $[\nabla^2 U \cdot \nabla U] := \sum_\alpha \nabla^2 U_\alpha \cdot \nabla U_\alpha$ . We include also many stored energies of Nonlinear Elasticity and many functionals of interest in Optimization, for which the lack of LSC requires relaxation methods.

We derive our MPs by constructing suitable energy comparison functions which satisfy (MP-v) and contradict minimality. Generally, the projection onto  $\overline{\text{co}}(U(\partial\Omega))$  would give a favorable deformation ([B-F2]). However, it is clear that in the absence of quasiconvexity we can not perform it to decrease  $E_\alpha$ . For this, we introduce a nonconvex method, the “Folding”. We obtain the desired deformations as weak limits of isometries which converge to the projection. The lack of weak\* LSC of  $E_\alpha$  is compensated by the combined use of the invariance (HR) and a suitable weak\* approximation of the projection onto convex sets by reflections. (HR) “compactifies”  $E_\alpha$ , which perceives the convergence as if it was strong.

The systematic study of  $(E_\infty)$  and (A) was initiated by Aronsson in [Ar1], [Ar2], [Ar4], [Ar5], [Ar6] who studied  $(E_\infty)$  with  $\mathcal{L}(\nabla u, u, x) = |\nabla u|$ ,  $N = 1, n = 2$ . Utilizing  $\|\cdot\|_{[L^p(\Omega)]^n}$  approximate problems to  $E_\infty(u) = \|\nabla u\|_{[L^\infty(\Omega)]^n}$ , in the limit of  $\Delta_p u = \text{Div}(|\nabla u|^{p-2} \nabla u)$  as  $p \rightarrow \infty$ , he derived the  $\infty$ -Laplacian:

$$\Delta_\infty u := (\nabla u)^\top [\nabla^2 u \cdot \nabla u] = \sum_{i,j} u_{x_i x_j} u_{x_j} u_{x_i} = 0. \quad (\Delta_\infty)$$

When  $\mathcal{L}(\nabla u, u, x) = |\nabla u|$ ,  $\infty$ -Harmonic functions relate to **AM**'s of the Lipschitz functional

$$\text{Lip}[U, \Omega] := \sup \left\{ \frac{|U(x) - U(y)|}{|x - y|} : x \neq y, x, y \in \Omega \right\}. \quad (\text{Lip})$$

For relevant material, we refer to Aronsson [Ar3], Aronsson-Crandall-Juutinen [A-C-J], Crandall [Cr1] and Gariépy-Wang-Yu [G-W-Y]. For recent applications of **(A)** see Barron-Evans-Jensen [B-E-J].

Our MP's for **(E<sub>1</sub>)** apply to Sobolev and for **(E<sub>∞</sub>)** to Lipschitz functions. The space setup is flexible enough to permit *anisotropic* exponents. It has been around in the literature for quite a long, but mostly in different guises. We also conjecture that **(MP-v)** can be derived for certain non-variational PDEs.

This paper is organized as follows. This Introduction is followed by Section 2, into which we fix our setup and introduce some basics. In Section 3 we present our assumptions and the main results. In Section 4 we present the auxiliary results and describe our methodology. Finally, we have collected together all of our proofs in Section 5.

**Acknowledgement.** I thank N. Alikakos for our discussions and his constructive criticism which lead to a more readable version of this manuscript. I also thank Ch. Sourdis and A. Quaas for communicating me some of my references and G. Paschalides for his advice on  $\text{\LaTeX}$ .

## 2 Basics.

**The functional space setup.** Let  $\Omega \subseteq \mathbb{R}^n$  be an open set and  $U = (U_1, \dots, U_N) : \Omega \rightarrow \mathbb{R}^N \equiv \mathbb{R}^{1 \times N}$  a Lebesgue measurable function,  $n, N \geq 1$ . The weak  $\nabla U$  when it exists is understood as a matrix function  $\nabla U = [\nabla U_1 | \dots | \nabla U_N] : \Omega \rightarrow \mathbb{R}^{n \times N}$ . The  $k$ -dimensional Lebesgue and Hausdorff measures are denoted by  $\mathcal{L}^k$ ,  $\mathcal{H}^k$  and the  $L^p$  and  $W^{1,p}$  spaces of vector functions by  $[L^p(\Omega)]^N$ ,  $[L^p_{\text{loc}}(\Omega)]^N$  and  $[W^{1,p}(\Omega)]^N$ ,  $[W^{1,p}_{\text{loc}}(\Omega)]^N$ ,  $[W^{1,p}_0(\Omega)]^N$ , etc. Let  $p, q \in [1, \infty]$ . We consider the  $(p, q)$ -Anisotropic Sobolev space:

$$[W^{1;p,q}(\Omega)]^N := \left\{ U \in [L^p(\Omega)]^N : \nabla U \in [L^q(\Omega)]^{n \times N} \right\} \quad (\text{AS})$$

(with a slight abuse of standard terminology on “anisotropies”). Obviously, we have  $[W^{1;p,p}(\Omega)]^N = [W^{1,p}(\Omega)]^N$ . The respective local space  $[W^{1;p,q}_{\text{loc}}(\Omega)]^N$  is defined analogously. **(AS)** becomes a Banach space, when equipped with the norm  $\|U\|_{[W^{1;p,q}(\Omega)]^N} := \|U\|_{[L^p(\Omega)]^N} + \|\nabla U\|_{[L^q(\Omega)]^{n \times N}}$ . Let now  $\mathcal{T}^w_{1;p,q}$  be a weak topology on  $[W^{1;p,q}(\Omega)]^N$ , depending on  $p, q$ . The *trace-0 subspace* of **(AS)** is the weak closure

$$[W^{1;p,q}_0(\Omega)]^N := \overline{[C^\infty_c(\Omega)]^N}^{\mathcal{T}^w_{1;p,q}}. \quad (\text{AfS})$$

The topology  $\mathcal{T}^w_{1;p,q}$  can be defined via the isometry  $[W^{1;p,q}(\Omega)]^N \hookrightarrow [L^p(\Omega)]^N \times [L^q(\Omega)]^{n \times N} : U \mapsto (U, \nabla U)$ . We take the weak topology of  $L^p \times L^q$ , which is weak-weak when  $1 \leq p, q < \infty$ , with weak\* instead of weak when  $p$  or  $q$  are  $\infty$ . Given  $U_0 \in [W^{1;p,q}(\Omega)]^N$ , we set  $[W^{1;p,q}_{U_0}(\Omega)]^N := U_0 + [W^{1;p,q}_0(\Omega)]^N$ .

**Weak formulation of  $U(\Omega) \subseteq \overline{\text{co}}(U(\partial\Omega))$ .** We introduce the necessary machinery in order to give a meaning to **(MP-v)** when  $U$  and  $\Omega$  are merely measurable. Let  $\mu$  be a positive Borel measure on  $\Omega$ ,  $U : \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^N$  a  $\mu$ -measurable function and  $A \subseteq \Omega$  with  $\mu(A) > 0$ . Let also “ $\mathbf{L}$ ” denote the restriction  $(\mu \mathbf{L} A)(E) := \mu(A \cap E)$ . We notice that the first definition (**Essmg**) is known.

**Definition.** The  $\mu\mathbf{L}A$ -essential range of  $U$  is defined as the closed set

$$\mu - \text{ess } U(A) := \left\{ u \in \mathbb{R}^N : \forall \varepsilon > 0, \mu(A \cap \{|U - u| < \varepsilon\}) > 0 \right\}. \quad (\text{Essrng})$$

Hence, we have  $\mu - \text{ess } U(A) = \left\{ u \in \mathbb{R}^N : \mu\mathbf{L}A - \text{ess } \inf |U - u| = 0 \right\}$ .  $\mu$  will be dropped if  $\mu = \mathcal{L}^n \mathbf{L}A$ .

**Definition.** We define the  $\mu\mathbf{L}A$ -essential closed convex hull of  $U$  as the set

$$\mu - \text{ess } \overline{\text{co}}(U(A)) := \bigcap \left\{ C : C \subseteq \mathbb{R}^N \text{ closed convex with } \mu(A \cap \{U \notin C\}) = 0 \right\}. \quad (\text{EssCo})$$

Let now  $\mu = \mathcal{L}^n$ . We extend  $\min_{\partial\Omega}$  and  $\max_{\partial\Omega}$  to  $N > 1$  by extending **(EssCo)** to lower dimensional subsets of  $\overline{\Omega}$  (cf. Puzzi-Serrin **[P-S]**). Let  $K \subseteq \overline{\Omega}$  be a Borel set with  $\text{dist}(u, K) := \inf \{|u - z| : z \in K\}$ .

**Definition.** The approximate closed convex hull of  $U$  over  $K$  is the closed set  $C_K \subseteq \mathbb{R}^N$  defined by:

$$\text{app } \overline{\text{co}}(U(K)) = C_K \Leftrightarrow \begin{cases} \text{for all } \varepsilon > 0, \text{ there is an open set } A_\varepsilon \subseteq \mathbb{R}^n \text{ with } K \subseteq A_\varepsilon \text{ such that} \\ C_K \subseteq \text{ess } \overline{\text{co}}(U(A_\varepsilon \cap \Omega)) \subseteq \{\text{dist}(\cdot, C_K) \leq \varepsilon\}. \end{cases} \quad (\text{Co})$$

In this case, we can compute  $C_K$  by means of:  $\text{app } \overline{\text{co}}(U(K)) = \liminf_{\varepsilon \rightarrow 0} \text{ess } \overline{\text{co}}(U(A_\varepsilon \cap \Omega))$ .

**(Co)** makes sense for merely measurable even unbounded  $U$ . It applies in particular to  $K = \partial\Omega$ . When there is a neighborhood  $A \supseteq K$  such that  $U \in [L^\infty(\Omega \cap A)]^N$ , an equivalent more flexible definition is

$$\text{app } \overline{\text{co}}(U(K)) := \lim_{\varepsilon \rightarrow 0^+} \text{ess } \overline{\text{co}}\left(U(\{\text{dist}(\cdot, K) < \varepsilon\} \cap \Omega)\right). \quad (\text{CO})$$

In the same spirit, the *approximate range of  $U$  over  $K$*  can be defined by

$$\text{app}(U(K)) := \lim_{\varepsilon \rightarrow 0^+} \text{ess}(U(\{\text{dist}(\cdot, K) < \varepsilon\} \cap \Omega)). \quad (\text{AppRng})$$

When  $N = 1$ , then **(Co)** reduces to  $\text{app } \overline{\text{co}}(u(\partial\Omega)) = [\min_{\partial\Omega} u, \max_{\partial\Omega} u]$ , with  $\max, \min$  understood as approximate liminf and limsup respectively (Evans-Gariepy **[E-G]**). When  $U \in [C^0(\Omega)]^N$ , we may set  $\overline{\text{co}}(U(K)) := \text{app } \overline{\text{co}}(U(K))$  keeping in mind that if  $\text{diam}(K) = \infty$ , this is understood as the limit **(CO)**.

Our convergence technique which applies when  $E_\alpha$  fails to be weakly\* LSC is based on the invariance of  $\mathcal{L}(\cdot, \xi, x)$  with respect to a set of reflections, which we now introduce.

**The set of  $\mathbb{Z}_2$ -reflection groups generated by a convex set  $C \subseteq \mathbb{R}^N$ .** The group  $\text{AO}(N, \mathbb{R})$  of *Affine Orthogonal Transformations* of  $\mathbb{R}^N$  is the set  $\mathbb{R}^N \times \text{O}(N, \mathbb{R})$ , equipped with the product  $(\xi, A) \cdot (\eta, B) := (A\eta + \xi, AB)$  and zero & inverse elements  $(0, I)$ ,  $(\xi, A)^{-1} = (-A^{-1}\xi, A^{-1})$ .  $\text{AO}(N, \mathbb{R})$  is the *Semidirect Product*  $\mathbb{R}^N \ltimes \text{O}(N, \mathbb{R})$ . Let  $\xi \in \mathbb{S}^{N-1}$  be a unit vector.  $\xi$  uniquely defines a hyperplane  $H_\xi := (\text{span}[\xi])^\perp \subseteq \mathbb{R}^N$  and a reflection  $R_\xi \in \text{O}(N, \mathbb{R})$  with respect to (w.r.t.)  $H_\xi$ , given by  $R_\xi u := u - 2(u \cdot \xi)\xi$ . If  $u_0 \in \mathbb{R}^N$ , the affine reflection w.r.t.  $H_\xi + u_0$  is given by  $u \mapsto u - 2((u - u_0) \cdot \xi)\xi = R_\xi u + 2(u_0 \cdot \xi)\xi$ . Let  $C \subseteq \mathbb{R}^N$  be closed and convex. The set of *Supporting Hyperplanes*  $\mathcal{S}(C)$  is

$$\mathcal{S}(C) = \left\{ H_\xi + u \in \mathbb{R}^N \mid \forall u \in \partial C, \exists \xi \in \mathbb{S}^{N-1} : \forall v \in C, \xi \cdot (u - v) \geq 0 \right\}. \quad (\mathcal{S}(C))$$

We will write  $\xi = \hat{u}$  for the normal to the hyperplane at  $u$ , but unless  $\partial C$  is  $C^1$ ,  $u \mapsto \hat{u}$  is multivalued. For any  $u \in \partial C$ ,  $\hat{u}$  generates an reflection  $w \mapsto R_{\hat{u}} w + 2(u \cdot \hat{u})\hat{u}$  in  $\text{AO}(N, \mathbb{R})$  w.r.t.  $H_{\hat{u}} + u$  and the induced  $R_{\hat{u}}$  in  $\text{O}(N, \mathbb{R}) \cap \{\det = -1\}$ . Since  $R_{\hat{u}}^2 = I$ , all  $\hat{u}$  generate cyclic groups  $\{I, R_{\hat{u}}\} =: [R_{\hat{u}}]$  isomorphic to  $\mathbb{Z}_2$ .

**Definition.** The set of  $\mathbb{Z}_2$ -reflection groups generated by (the supporting hyperplanes of)  $C$  is:

$$\mathcal{R}(C) := \{[R_{\hat{u}}] : u \in \partial C\} = \{I\} \cup \{R_{\hat{u}} : u \in \partial C\}. \quad (\mathcal{R}(C))$$

### 3 Assumptions and Main Results.

**Hypotheses on  $\mathcal{L}$  and the Folding Principles.** Let  $\mathcal{L} : \mathbb{R}^{n \times N} \times \mathbb{R}^N \times \Omega \rightarrow [0, \infty)$  be a Carathéodory function,  $\Omega \subseteq \mathbb{R}^n$  open set. We shall need to impose the following hypotheses:

$$\left\{ \begin{array}{l} \text{For all } P \in \mathbb{R}^{n \times N} \text{ and } \mathcal{L}^n\text{-}\Omega\text{-a.e. } x, \text{ we assume that } \mathcal{L}(P, \cdot, x) \in C^{0+\gamma}(\mathbb{R}^N) \\ \text{for some } \gamma \in (0, 1], \text{ locally uniformly in } (P, x) : \text{ For any } B' \times B \times \Omega' \subset \subset \\ \mathbb{R}^{n \times N} \times \mathbb{R}^N \times \Omega \text{ and all } (P, \bar{\xi}, x), (P, \xi, x) \in B' \times B \times \Omega', \\ \left| \mathcal{L}(P, \bar{\xi}, x) - \mathcal{L}(P, \xi, x) \right| \leq C(B', \Omega'; B) \left| \bar{\xi} - \xi \right|^\gamma \\ \text{with} \\ C(B', \Omega'; B) := \sup_{P \in B'} \operatorname{ess\,sup}_{x \in \Omega'} \left( [\mathcal{L}(P, \cdot, x)]_{C^{0+\gamma}(B)} \right) < \infty. \end{array} \right. \quad (\text{HL})$$

$$\left\{ \begin{array}{l} \text{There is a compact convex set } C \subseteq \mathbb{R}^N \text{ such that for } \mathcal{L}^n\text{-}\Omega\text{-a.e. } x \text{ and all } \xi \\ \text{in } \mathbb{R}^N, \mathcal{L}(\cdot, \xi, x) \text{ is invariant under the action of the set } \mathcal{R}(C) \text{ of reflection} \\ \mathbb{Z}_2 \text{ groups generated by } \mathcal{S}(C): \\ \mathcal{L}(PR, \cdot, \cdot) = \mathcal{L}(P, \cdot, \cdot), \\ \text{for all } P \in \mathbb{R}^{n \times N} \text{ and all } R \in \mathcal{R}(C) \subseteq \text{O}(N, \mathbb{R}) \cap \{\det = -1\}. \end{array} \right. \quad (\text{HR})$$

$$\left\{ \begin{array}{l} \text{There is a compact convex set } C \subseteq \mathbb{R}^N \text{ such that for } \mathcal{L}^n\text{-}\Omega\text{-a.e. } x \text{ and all } P \\ \text{in } \mathbb{R}^{n \times N}, \\ \left\{ \mathcal{L}(P, \cdot, x) \leq \sup_{\bar{\xi} \in \partial C} \mathcal{L}(P, \bar{\xi}, x) \right\} \subseteq C. \end{array} \right. \quad (\text{HC})$$

These assumptions suffice to derive our MPs for  $(E_\infty)$ . Concerning  $(E_1)$ , we need two more hypotheses.

$$\left\{ \begin{array}{l} \text{We assume that (HL) holds uniformly in all } (P, x) \in \mathbb{R}^{n \times N} \times \Omega \text{ for } \gamma = 1. \\ \text{That is, } \mathcal{L}(P, \cdot, x) \in W_{\text{loc}}^{1,\infty}(\mathbb{R}^N) \text{ and for any } B \subset \subset \mathbb{R}^N, \\ \sup_{P \in \mathbb{R}^{n \times N}} \operatorname{ess\,sup}_{x \in \Omega} \left( \operatorname{Lip} [\mathcal{L}(P, \cdot, x), B] \right) < \infty. \end{array} \right. \quad (\text{HL1})$$

$$\left\{ \begin{array}{l} \text{There exist } q \geq 1, C > 0 \text{ and a continuous } \beta \in C^0(\mathbb{R}^N \times \Omega) \text{ with } \beta(\xi, \cdot) \\ \text{in } L^1(\Omega) \text{ for all } \xi \in \mathbb{R}^N, \text{ such that} \\ \mathcal{L}(P, \xi, x) \leq C |P|^q + \beta(\xi, x). \end{array} \right. \quad (\text{HB})$$

Evidently, (HC) is equivalent to the existence of a  $C \subseteq \mathbb{R}^N$  compact convex: for all  $\xi \in \mathbb{R}^N \setminus C$  and all  $\bar{\xi} \in \partial C$ ,  $\mathcal{L}(\cdot, \xi, \cdot) > \mathcal{L}(\cdot, \bar{\xi}, \cdot)$ . Let e.g.  $\mathcal{L}$  be of the frequent decoupled form  $\mathcal{L}(P, \xi, x) := F(P, x) + h(x)G(\xi)$ ,  $h \geq 0$ . Then, (HC) means that  $\mathbb{R}^N \setminus C$  is contained into a suplevel set  $\{G > t\}$  of  $G$ . We now present the MPs for  $(E_1)$  and the PDE (E). Recall the definitions (Essrng), (EssCo), (Co), (CO).

**Theorem 3.1. (Folding Principle for  $(E_1)$  (I))** Let  $\Omega \subseteq \mathbb{R}^n$  be an open domain with  $\mathcal{L}^n(\Omega) < \infty$  and  $\mathcal{L} : \mathbb{R}^{n \times N} \times \mathbb{R}^N \times \Omega \rightarrow [0, \infty)$  a Carathéodory function satisfying (HL), (HL1) and (HB). Let also  $C$

be a compact convex set with respect to which  $\mathcal{L}$  satisfies (HR) and (HC). If  $U_0 \in [W^{1;p,q}(\Omega)]^N$  with  $p \in [1, \infty]$  and  $q$  as in (HB), satisfying

$$\text{app } \overline{\text{co}}(U_0(\partial\Omega)) \subseteq C, \quad (\text{BC})$$

then, every minimizing sequence  $(U_i)_1^\infty$  of the functional  $(E_1)$  in  $[W_{U_0}^{1;p,q}(\Omega)]^N$ :

$$E_1(U_i) \longrightarrow \inf \left\{ E_1(U, \Omega) : U \in [W_{U_0}^{1;p,q}(\Omega)]^N \right\}, \quad \text{as } i \rightarrow \infty, \quad (\text{P1})$$

satisfies the a priori localization:

$$\bigcup_1^\infty \text{ess } U_i(\Omega) \subseteq C. \quad (\text{FP}_1 \text{ (I)})$$

Theorem 3.1 derives a weaker version of (MP-v), but in a general context. Even when a solution to (P1) ceases to exist, minimizing sequences are localized into  $C \subseteq \mathbb{R}^N$ , if their boundary values are inside  $C$ . In fact, (FP<sub>1</sub> (I)) is analogous to a weaker (MP-s) which reads  $\sup_\Omega[\pm u] \leq \max_{\partial\Omega}[\pm u] + C$  for  $C > 0$ .

Now we strengthen (FP<sub>1</sub> (I)). We refer to Dacorogna [D] for sufficient conditions under which a minimizing sequence of (P1) converges to a minimizer  $U^*$  of the relaxed functional:

$$QE_1(U, \Omega) := \int_\Omega Q\mathcal{L}(\nabla U, U, x) dx, \quad (QE_1)$$

where  $Q\mathcal{L}$  is the  $M$ -quasiconvex envelope of  $\mathcal{L}$ , defined by the right hand side of (M-QC). We recall that the relaxed solution  $U^*$  to (P1) (equivalently, the Young measure solution) generally satisfies

$$\begin{aligned} E_1(U_i, \Omega) &\longrightarrow QE_1(U^*, \Omega) = \inf \left\{ QE_1(U, \Omega) : U \in [W_{U_0}^{1;p,q}(\Omega)]^N \right\} \\ &= \inf \left\{ E_1(U, \Omega) : U \in [W_{U_0}^{1;p,q}(\Omega)]^N \right\} \end{aligned} \quad (\text{QP1})$$

and

$$\begin{cases} \text{Div}(\mathcal{L}_P(\nabla U_i, U_i, x)) - \mathcal{L}_\xi(\nabla U_i, U_i, x) \longrightarrow 0, & \text{as } i \rightarrow \infty \text{ in } ([W_{U_0}^{1;p,q}(\Omega)]^N)^*, \\ \text{Div}((Q\mathcal{L})_P(\nabla U^*, U^*, x)) - (Q\mathcal{L})_\xi(U^*, U^*, x) = 0. \end{cases}$$

**Theorem 3.2. (Folding Principle for  $(E_1)$  (II))** In the setting of Theorem 3.1, we additionally assume

$$U_i \rightharpoonup U^*, \quad \text{in the weak topology } \mathcal{T}_{1;p,q}^w \text{ of } [W_{U_0}^{1;p,q}(\Omega)]^N.$$

Then, the relaxed solution  $U^*$  to (P1) which is a solution to (QP1) satisfies:

$$\text{ess } U^*(\Omega) \subseteq C. \quad (\text{FP}_1 \text{ (II)})$$

Now we derive the stronger conclusion (MP-v).

**Corollary 3.3. (Folding Principle for  $(E_1)$  (III))** In the setting of Theorems 3.1 and 3.2, if  $U = U_i, U^*$  respectively, we additionally assume that given  $U_0$  in  $[W^{1;p,q}(\Omega)]^N$ ,

$$\mathcal{L} \text{ satisfies (HR) and (HC) with } C = \text{app } \overline{\text{co}}(U_0(\partial\Omega))$$



(which holds for all  $U_0$  when e.g.  $\mathcal{L}(P, \xi, x) = \mathcal{L}(P, x)$  with  $P \mapsto \mathcal{L}(P, \cdot)$  reflection invariant). Then:

$$\text{ess } U(\Omega) \subseteq \text{app } \overline{\text{co}} \left( U_0(\partial\Omega) \right). \quad (FP_1 \text{ (III)})$$

For general conditions guaranteeing the existence of minimizers to  $(E_1)$ , we refer to Dacorogna [D]. When our assumptions (HR) and (HC) are reduced to  $N = 1$ , their simpler scalar version reads

$$\text{Scalar } (N = 1) \text{ (HR)} : \quad \left\{ \begin{array}{l} \text{For all } P \in \mathbb{R}^n, \quad \mathcal{L}(-P, \cdot, \cdot) = \mathcal{L}(P, \cdot, \cdot). \end{array} \right. \quad (\text{HR1})$$

$$\text{Scalar } (N = 1) \text{ (HC)} : \quad \left\{ \begin{array}{l} \text{There exist } \xi_{\min} < \xi_{\max} \text{ in } \mathbb{R} : \text{ for all } \xi < \xi_{\min} \text{ and } \xi > \xi_{\max}, \\ \mathcal{L}(\cdot, \xi, \cdot) > \max \left\{ \mathcal{L}(\cdot, \xi_{\min}, \cdot), \mathcal{L}(\cdot, \xi_{\max}, \cdot) \right\}. \end{array} \right. \quad (\text{HC1})$$

**Corollary 3.4. (Folding Principle for the PDE (E))** Let  $\mathcal{L}$ ,  $\Omega$ ,  $C$ ,  $p$ ,  $q$  be as in Theorem 3.1 with  $\mathcal{L}(\cdot, \cdot, x) \in C^1(\mathbb{R}^{n \times N} \times \mathbb{R}^N)$  for  $\mathcal{L}^n \text{--}\Omega$ -a.e.  $x$ . If  $N \geq 1$ , minimizing weak solutions  $U \in [W^{1;p,q}(\Omega)]^N$  to (E) satisfy

$$\left\{ \begin{array}{ll} \text{ess } U(\Omega) \subseteq C, & \text{if } \text{app } \overline{\text{co}} \left( U(\partial\Omega) \right) \subseteq C, \\ \text{ess } U(\Omega) \subseteq \text{app } \overline{\text{co}} \left( U(\partial\Omega) \right), & \text{if } C = \text{app } \overline{\text{co}} \left( U(\partial\Omega) \right) \text{ in (HR), (HC)}. \end{array} \right.$$

When  $N = 1$ , this reduces for scalar minimizing solutions  $u \in W^{1;p,q}(\Omega)$  of (E) respectively to:

$$\left\{ \begin{array}{l} \xi_{\min} \leq \inf_{\Omega} u \leq \sup_{\Omega} u \leq \xi_{\max}, \\ \min_{\partial\Omega} u \leq \inf_{\Omega} u \leq \sup_{\Omega} u \leq \max_{\partial\Omega} u. \end{array} \right.$$

We now apply Corollary 3.4 to the A-Laplacian (Puzzi-Serrin [P-S]). We have not assumed (M-QC), so 3.4 holds for non-elliptic (E). Our only growth bound (HB) includes the Minimal Surface operator.

**Corollary 3.5. (Folding Principle for Vectorial A-Harmonic Functions)** If  $\mathcal{L}$ ,  $\Omega$ ,  $p$ ,  $q$  are as in Theorem 3.1 with  $\mathcal{L} = \mathcal{L}(P, x)$  and for  $\mathcal{L}^n \text{--}\Omega$ -a.e.,  $\mathcal{L}(\cdot, x)$  is reflection invariant and in  $C^1(\mathbb{R}^{n \times N})$ , then minimizing vectorial weak solutions  $U \in [W^{1;p,q}(\Omega)]^N$  to  $\text{Div}(\mathcal{L}_P(\nabla U, x)) = 0$  can be chosen to satisfy for any  $\Omega' \subseteq \Omega$ :

$$\text{ess } U(\Omega') \subseteq \text{app } \overline{\text{co}} \left( U(\partial\Omega') \right).$$

In particular, for  $\mathcal{L}(\nabla U, x) := \int_0^{|\nabla U|} sA(s) ds$ , A-Harmonic functions which solve  $\text{Div}(A(|\nabla U|)\nabla U) = 0$  can be chosen to satisfy it. The latter holds for all  $U$  in the convex case.

**Application 3.6. (Hyperelastic materials tend to take up convex portions of the space)** We exhibit an application of Theorem 3.1 to Elasticity. Let  $\Omega \subseteq \mathbb{R}^3$  be the reference configuration of a hyperelastic material  $\mathcal{M}$  and  $\mathcal{L} : \mathbb{R}^{3 \times 3} \times \Omega \rightarrow \mathbb{R}$  the stored energy. We assume  $\mathcal{L}(P, x) = W(|P|, x)$ ,  $W$  in  $C^0([0, \infty) \times \Omega)$ . Hence  $\mathcal{L}$  is objective and isotropic (Dacorogna [D]). We look for a deformation  $U$  in  $[W_{U_0}^{1;p}(\Omega)]^N$ , solving (E) with  $\det(\nabla U) > 0$ . If phase transitions occur,  $U$  may not exist. Then, minimizing sequences  $(U_i)_1^\infty$  are finer and finer mixtures of different phases. By Theorem 3.1, there is  $(U_i)_1^\infty : \bigcup_1^\infty \text{ess } U_i(\Omega) \subseteq \text{app } \overline{\text{co}} \left( U_0(\partial\Omega) \right)$ . If  $\text{app } \left( U_0(\partial\Omega) \right)$  is convex,  $\mathcal{M}$  tends in average to take up a convex portion of  $\mathbb{R}^3$ .



Now we turn to the MPs for  $(E_\infty)$  and  $(A)$ , choosing  $p = q = \infty$  in  $(AS)$ . Sufficient conditions for weak\*-LSC of  $(E_\infty)$  and for the existence of  $AM$ 's appear in Barron-Jensen-Wang [BJW1], [BJW2]. Due to the properties of  $[W^{1,\infty}(\Omega)]^N$ , we may drop “ $\mathcal{L}^n(\Omega) < \infty$ ” and the assumptions  $(HL1)$ ,  $(HB)$ . By continuity, the definitions  $(EssCo)$ ,  $(Co)$  and  $(CO)$  now have a classical meaning, at least when  $\text{diam}(\Omega) < \infty$ .

**Theorem 3.7. (Folding Principle for  $(E_\infty)$  (I))** Let  $\mathcal{L} : \mathbb{R}^{n \times N} \times \mathbb{R}^N \times \Omega \longrightarrow [0, \infty)$  be a Carathéodory function,  $\Omega \subseteq \mathbb{R}^n$  open and  $C \subseteq \mathbb{R}^N$  compact convex. We assume that  $\mathcal{L}$  satisfies  $(HL)$ , together with  $(HR)$  and  $(HC)$  for this  $C$ . If  $U_0 \in [W^{1,\infty}(\Omega)]^N$ , with

$$\overline{co}(U_0(\partial\Omega)) \subseteq C,$$

then, every absolute minimizer  $U^* \in [W^{1,\infty}_{U_0}(\Omega)]^N \cap AM(E_\infty(-, \Omega))$  of the functional  $(E_\infty)$ , satisfies:

$$U^*(\Omega) \subseteq C. \quad (FP_\infty \text{ (I)})$$

The stronger conclusion  $(MP-v)$  is also in order, exactly as in the integral case of  $(E_1)$ .

**Corollary 3.8. (Folding Principle for  $(E_\infty)$  (II))** Let  $\mathcal{L}$ ,  $\Omega$ ,  $C$  be as in Theorem 3.7 with  $U^* \in [W^{1,\infty}(\Omega)]^N \cap AM(E_\infty(-, \Omega))$ . If

$$\mathcal{L} \text{ satisfies } (HR) \text{ and } (HC) \text{ with } C = \overline{co}(U^*(\partial\Omega))$$

(which holds always when e.g.  $\mathcal{L}(P, \xi, x) = \mathcal{L}(P, x)$  with  $P \mapsto \mathcal{L}(P, \cdot)$  reflection invariant), then:

$$U^*(\Omega) \subseteq \overline{co}(U^*(\partial\Omega)). \quad (FP_\infty \text{ (II)})$$

We refer to Barron-Jensen-Wang [BJW1], Crandall [Cr2] and Crandall-Wang-Yu [C-W-Y] for sufficient conditions under which  $AM$ 's are viscosity solutions of  $(A)$ . For regular solutions when  $N > 1$ , consult Barron-Jensen-Wang [BJW2]. When  $N = 1$ , we recall the scalar assumption  $(HC1)$  and  $(HR1)$ .

**Corollary 3.9. (Folding Principle for the PDE  $(A)$ )** In the setting of Theorem 3.7 and Corollary 3.8, we additionally assume  $\mathcal{L} \in C^1(\mathbb{R}^{n \times N} \times \mathbb{R}^N \times \Omega)$ .

(i) When  $N = 1$ , absolutely minimizing viscosity solutions  $u^* \in W^{1,\infty}(\Omega)$  to the scalar PDE  $(A)$

$$\mathcal{L}_P(\nabla u, u, x) \left( \nabla^2 u \left( \mathcal{L}_P(\nabla u, u, x) \right)^\top + \mathcal{L}_\xi(\nabla u, u, x) \nabla u + \mathcal{L}_x(\nabla u, u, x) \right) = 0$$

satisfy in the case of Theorem 3.7 and Corollary 3.8 respectively

$$\begin{cases} \xi_{\min} \leq \inf_{\Omega} u^* \leq \sup_{\Omega} u^* \leq \xi_{\max}, \\ \min_{\partial\Omega} u^* \leq \inf_{\Omega} u^* \leq \sup_{\Omega} u^* \leq \max_{\partial\Omega} u^*. \end{cases}$$

(ii) When  $N > 1$ , vectorial  $AM$  regular solutions  $U^*$  to the PDE system  $(A)$  satisfy respectively

$$\begin{cases} U^*(\Omega) \subseteq C, \\ U^*(\Omega) \subseteq \overline{co}(U^*(\partial\Omega)). \end{cases}$$

We now consider the “extremal” members of the  $p$ -Harmonic functions as  $p \rightarrow \infty$ , the  $\infty$ -Harmonic.

**Corollary 3.10. (Folding Principle for the generalized  $\infty$ -Laplace system)** *If  $\mathcal{L} = \mathcal{L}(P, x)$  with  $\mathcal{L} \in C^1(\mathbb{R}^{n \times N} \times \Omega)$  and  $P \mapsto \mathcal{L}(P, \cdot)$  reflection invariant, regular **AM** solutions  $U^*$  to the generalized  $\infty$ -Laplace system of PDEs*

$$\mathcal{L}_P(\nabla U, x) \left( \left[ \nabla^2 U \cdot (\mathcal{L}_P(\nabla U, x))^\top \right] + \mathcal{L}_x(\nabla U, x) \right) = 0,$$

*can be chosen to satisfy for any subdomain  $\Omega' \subseteq \Omega$ :*

$$U^*(\Omega') \subseteq \overline{\text{co}} \left( U^*(\partial\Omega') \right).$$

*If  $\mathcal{L}(\nabla U, x) := |\nabla U|$ , this holds for all **AM**  $\infty$ -Harmonic functions, solutions to  $(\nabla U)^\top [\nabla^2 U \cdot \nabla U] = 0$ .*

Corollaries 3.9, 3.10 extends the Maximum Principle of Jensen [Je2] to more general  $\mathcal{L}$  and to  $N > 1$ .

**Remark 3.11.** *All of our MPs admit sharper versions at the expense of additional complications, obtained by imposing a constraint, relaxing the boundedness of  $C$ , and so on. We refrain from this task, since only technical modifications are required. Also, our technique seems to extend directly to the  $[BV(\Omega)]^N$  setting for the relaxed version of  $(E_\alpha)$  of linear growth, utilizing the chain rule formula of Ambrosio-Dal Maso (see [A-F-P]), thus extending the results of Bildhauer-Fuchs [B-F2].*

## 4 Auxiliary Results and Methodology.

The set  $\mathcal{R}(C)$  involved in our assumption (HR) plays a crucial role for us. As such, we wish to examine its structure. For compact  $C$ ,  $\mathcal{R}(C)$  turns out to be a familiar manifold. We recall from Mattila [M] that a subset  $E \subseteq \mathbb{R}^M$  is  $m$ -rectifiable,  $m \in \{0, 1, \dots, M\}$ , if there exist  $(g_i)_1^\infty \subseteq [W^{1,\infty}(\mathbb{R}^m)]^M$  and  $(A_i)_1^\infty \subseteq \mathbb{R}^m$ :

$$\mathcal{H}^m \left( E \setminus \bigcup_1^\infty g_i(A_i) \right) = 0.$$

**Lemma 4.1. (Structure of  $\mathcal{R}(C)$ )** *Let  $C \subseteq \mathbb{R}^N$  be a given closed convex set with  $\text{int}(C) \neq \emptyset$ . Then:*

- (i) *The set  $\mathcal{R}(C)$  is an  $(N-1)$ -rectifiable subset of  $O(N, \mathbb{R}) \subseteq \mathbb{R}^{N^2}$ .*
- (ii)  *$\mathcal{R}(C) \setminus \{I\}$  is a path-connected subset of  $O(N, \mathbb{R}) \cap \{\det = -1\}$ .*
- (iii) *If  $C$  is compact,  $\mathcal{R}(C) \setminus \{I\}$  is a compact manifold of reflection matrices with  $\dim(\mathcal{R}(C)) = N-1$ , invariant within  $\{C \subseteq \mathbb{R}^N : C \text{ compact, convex and } \text{int}(C) \neq \emptyset\}$ . That is, if  $\mathbb{B}^N = \mathbb{B}_1^N(0)$  is the ball,*

$$\mathcal{R}(C) \setminus \{I\} = \mathcal{R}(\mathbb{B}^N) \setminus \{I\} \cong \mathbb{R}P^{N-1},$$

*where  $\mathbb{R}P^{N-1}$  is the projective space.*

- (iv) *In the case of non-compact  $C$ , the rectifiable set  $\mathcal{R}(C)$  has an injection into  $\mathcal{R}(\mathbb{B}^N)$ .*

**Description of the convergence technique.** Let  $U$  be a “minimal” function for  $(E_\alpha)$ . Roughly, in order to establish  $U(\Omega) \subseteq C$  (or (MP-v) if  $C = \overline{\text{co}}(U(\partial\Omega))$ ), we assume to the contrary  $U(\Omega) \not\subseteq C$ . The

projected  $\mathcal{P}^C(U)$  satisfies  $\mathcal{P}^C(U)(\Omega) \subseteq C$ , so if  $E_\alpha(\mathcal{P}^C(U), \Omega) < E_\alpha(U, \Omega)$ ,  $\mathcal{P}^C(U)$  would be a suitable comparison function. However, this may not hold if (quasi)convexity fails. We then *weakly\** approximate  $\mathcal{P}^C$  by Lipschitz maps  $\mathcal{F}_m^C$  composed of reflections with respect to the supporting hyperplanes of  $C$ :  $\mathcal{F}_m^C \xrightarrow{*} \mathcal{P}^C$  as  $m \rightarrow \infty$  in  $[W_{loc}^{1,\infty}(\mathbb{R}^N)]^N$  (see  $(\diamond)$ ). In each step, we fold by reflections an arbitrarily large ball, arbitrarily close to  $\partial C$ . We then prove that  $\mathcal{F}_m^C(U) \rightarrow \mathcal{P}^C(U)$  in  $[W^{1,p,q}(\Omega)]^N$ . By (HR), we have  $\mathcal{L}(\nabla(\mathcal{F}_m^C(U)), \mathcal{F}_m^C(U), \cdot) = \mathcal{L}(\nabla U, \mathcal{F}_m^C(U), \cdot)$ . Then, we show that  $\lim_{m \rightarrow \infty} E_\alpha(\mathcal{F}_m^C(U), \Omega)$  exists and

$$\begin{aligned} \lim_{m \rightarrow \infty} E_\alpha(\mathcal{F}_m^C(U), \Omega) &= \lim_{m \rightarrow \infty} \left\| \mathcal{L}(\nabla(\mathcal{F}_m^C(U)), \mathcal{F}_m^C(U), \cdot) \right\|_{L^\alpha(\Omega)} \\ &= \left\| \mathcal{L}(\nabla U, \mathcal{P}^C(U), \cdot) \right\|_{L^\alpha(\Omega)} \\ &< \left\| \mathcal{L}(\nabla U, U, \cdot) \right\|_{L^\alpha(\Omega)} \\ &= E_\alpha(U, \Omega). \end{aligned}$$

In the absence of (M-QC), (BJW-QC), the limit may not bound  $E_\alpha(\mathcal{P}^C(U), \Omega)$  although it exists and is realized as an  $\|\cdot\|_{L^\alpha(\Omega)}$ -norm, but it is not the  $E_\alpha$ -energy of a function. However, for any  $\varepsilon > 0$  small, exists  $m \in \mathbb{N}$ :  $\mathcal{F}_m^C(U)(\Omega) \subseteq \{\text{dist}(\cdot, C) < \varepsilon\}$ ,  $U(\Omega) \cap \{\text{dist}(\cdot, C) \geq \varepsilon\} \neq \emptyset$  and  $E_\alpha(\mathcal{F}_m^C(U), \Omega) < E_\alpha(U, \Omega)$ . Hence,  $U(\Omega) \subseteq \{\text{dist}(\cdot, C) < \varepsilon\}$ , for all  $\varepsilon > 0$ .

Roughly, this description follows our approach for  $(E_\infty)$ . For  $(E_1)$ , further complications occur due to the possible escape of mass to infinity because of the unboundedness of  $L^p$  functions.

We begin with a lemma which is the core of the process. We first need to supplement convexity with an auxiliary notion which rectifies the lack of smoothness. It is later dropped by approximation.

**Definition 4.2.** Let  $C$  be a simplex, i.e. a compact set  $C = \bigcap \{H_j^+ : 1 \leq j \leq K\}$  generated by halfspaces  $H_j^+$  determined by hyperplanes  $H_j = (\text{span}[\xi_j])^\perp + u_j$ . The reduced set  $\mathcal{R}^*(C) \subseteq \mathcal{R}(C)$  of  $\mathbb{Z}_2$ -reflection groups  $[R_{\tilde{u}}]$  generated by the sides  $H_j^* := H_j \cap C$  is the following discrete counterpart of  $\mathcal{R}(C)$ :

$$\mathcal{R}^* \left( \bigcap \{H_j^+ : 1 \leq j \leq K\} \right) := \{[R_{\xi_j}] : 1 \leq j \leq K\} = \{I\} \cup \{R_{\xi_j} : 1 \leq j \leq K\}. \quad (\mathcal{R}^*(C))$$

The algebraic span of  $\mathcal{R}^*(C)$  in  $O(N, \mathbb{R})$  is known as a *Coxeter group*, i.e. a finite reflection group.

**Definition 4.3. (Obtuseness)** (i) A simplex  $C \subseteq \mathbb{R}^N$  will be called obtuse, if its defining sides  $H_j^*$  meet in obtuse angles  $\geq \frac{\pi}{2}$ , that is, when for all  $i, j \in \{1, \dots, K\}$  we have the implication

$$H_i^* \cap H_j^* \neq \emptyset \Rightarrow 0 \leq \xi_i \cdot \xi_j \leq 1. \quad (\text{O})$$


(ii) A closed convex set  $C \subseteq \mathbb{R}^N$  will be called obtuse if it can be approximated from the outside by a decreasing sequence of obtuse simplices  $(C_m)_1^\infty$  whose sides support  $C$ :

$$C_m = \bigcap \{H_{m,j}^+ : 1 \leq j \leq K(m)\} \searrow C \quad \text{as } m \rightarrow \infty, \quad (\text{O1})$$

where  $H_{m,j} \in \mathcal{S}(C)$  with  $K(m) \geq N + 1$ , for all indices  $(m, j)$ .

(O1) is satisfied by smooth convex sets, obtuse simplices themselves (e.g. rectangles, cubes) and their intersections. However, “tear-drop” shaped sets and oblique simplices may not verify it (Fig. 1). Smooth

convex sets verify (O1) since the edges of any rectangle with sides supporting it are a positive distance away from it. Thus, the intersections of such rectangles define obtuse simplices.



(Fig. 1)

**Lemma 4.4. (Folding Map for Obtuse Convex Sets, (O1))** Let  $C \subseteq \mathbb{R}^N$  be a compact obtuse convex set verifying (O1) with  $0 \in \text{int}(C) \neq \emptyset$ . Let also  $A \subseteq \mathbb{R}^N$  be bounded.

For any  $\lambda > 1$ , there exists a non-expansive  $\mathcal{F}_\lambda^C : \mathbb{R}^N \rightarrow \mathbb{R}^N$  in  $[W_{loc}^{1,\infty}(\mathbb{R}^N)]^N$  with  $\text{Lip}[\mathcal{F}_\lambda^C] = 1$  and

$$\begin{cases} \mathcal{F}_\lambda^C(E \cap C) = E \cap C, & \text{for all } E \subseteq \mathbb{R}^N \\ \mathcal{F}_\lambda^C(A \setminus C) \subseteq \lambda C \setminus \text{int}(C). \end{cases} \quad (*)$$

$\mathcal{F}_\lambda^C$  consists of composites of affine reflections in  $\mathbb{R}^N \times \mathcal{R}(C) \subseteq \text{AO}(N, \mathbb{R})$  and depends on  $A$  only through  $a := \sup \{|u| : u \in A\}$ . If  $C$  is an obtuse simplex  $\cap \{H_j^+ : 1 \leq j \leq K\}$ , the reflections belong to  $\mathcal{R}^*(C)$ .

Hence, for every “nice” convex  $C$  and any bounded  $A$ , there is a finite sequence of successive reflections which leaves  $C$  invariant and folds  $A \setminus C$  as close to  $\partial C$  as desired.  $\mathcal{F}_\lambda^C$  will be applied to  $A = U(\Omega)$ , for  $U : \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^N$ . Our passing to the limit as  $\lambda \rightarrow 1^+$  gives rise to the projection  $\mathcal{P}^C : \mathbb{R}^N \rightarrow C$ .

**Lemma 4.5. (The Projection onto Convex Sets as Limit of Folding Maps)** Let  $C \subseteq \mathbb{R}^N$  be compact convex with  $0 \in \text{int}(C)$ . The family  $\{\mathcal{F}_\lambda^C : \lambda > 1\}$  of Lemma 4.4 is sequentially weakly\* precompact in  $[W_{loc}^{1,\infty}(\mathbb{R}^N)]^N$  and the Projection  $\mathcal{P}^C : \mathbb{R}^N \rightarrow C$  onto  $C$  is an accumulation point as  $\lambda \rightarrow 1^+$ .

We may now proceed to the deformation of functions  $U : \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^N$  via foldings. In order to treat essentially unbounded functions, we introduce the vector truncates: if  $U$  is  $\mathcal{L}^n \llcorner \Omega$ -measurable, we set

$$U^m := \begin{cases} U, & \text{on } \{|U| \leq m\} \subseteq \Omega \\ \frac{mU}{|U|}, & \text{on } \{|U| > m\} \subseteq \Omega, \quad m = 1, 2, \dots \end{cases} \quad (T)$$

We have  $\|U^m\|_{[L^\infty(\Omega)]^N} \leq m$ . The functions  $\mathcal{F}_m$  appearing in the next Lemma 4.6 are obtained by a double approximation argument (see  $(\diamond)$ ). In addition, the failure of reflexivity of  $L^1$  requires the use of measures. Let  $[\mathcal{M}(\Omega)]^{n \times N}$  be the space of  $\mathbb{R}^{n \times N}$ -valued Radon measures, equipped with the Fréchet topology induced by the seminorms  $\|\mu\|(\Omega') = \sup \left\{ \int_{\Omega'} \phi : d\mu \mid \phi \in [C_c^1(\Omega')]^{n \times N}, |\phi| \leq 1 \right\}$ ,  $\Omega' \subset\subset \Omega$ , with  $A : B = \text{tr}(AB^T)$ . Given a real  $\mu \in \mathcal{M}(\Omega)$  and  $f : \Omega \rightarrow \mathbb{R}^k$   $\mu$ -measurable, following Evans-Gariepy [E-G], the restriction  $\mu \llcorner f$  can be defined as  $(\mu \llcorner f)(A) := \int_A f d\mu$ , with  $\mu \llcorner f \in [\mathcal{M}(\Omega)]^k$ . If  $f = \chi_E$ , the characteristic of  $E \subseteq \mathbb{R}^n$ , it coincides with the restriction on  $E$ :  $(\mu \llcorner \chi_E)(A) = (\mu \llcorner E)(A) = \mu(E \cap A)$ .

**Lemma 4.6. (Convergence of Folding Deformations)** Let  $C \subseteq \mathbb{R}^N$  be compact convex with  $0 \in \text{int}(C)$ ,  $\Omega \subseteq \mathbb{R}^n$  open,  $\mathcal{P}^C : \mathbb{R}^N \rightarrow C$  the projection and  $\mathcal{F}_m^C : \mathbb{R}^N \rightarrow \mathbb{R}^N$  the Folding Maps of  $(\diamond)$ ,  $m \in \mathbb{N}$ .

(i) If  $U \in [L^p(\Omega)]^N$ , then up to suitable subsequences of the truncates  $(\mathbf{T})$  of  $U$  as  $m \rightarrow \infty$ ,

$$\mathcal{F}_m^C(U^m) \rightarrow \mathcal{P}^C(U) \quad \text{in } [L^p(\Omega)]^N, \quad p \in [1, \infty] \quad \text{and } \mathcal{L}^n \llcorner \Omega - \text{a.e.} \quad (\text{C1})$$

If  $\mathcal{L}^n(\Omega) = \infty > p$ , we additionally assume that  $\mathcal{L}^n(\{x \notin C\}) = \mathcal{L}^n(\{x \in \Omega : U(x) \notin C\}) < \infty$ .

(ii) If in addition  $U \in [W_0^{1;p,q}(\Omega)]^N$ , then up to suitable subsequences as  $m \rightarrow \infty$


$$\nabla(\mathcal{F}_m^C(U^m)) \xrightarrow{*} \nabla(\mathcal{P}^C(U)) \quad \text{in } [L^\infty(\Omega)]^{n \times N}, \quad \text{if } q = \infty, \quad (\text{C2a})$$

$$\nabla(\mathcal{F}_m^C(U^m)) \rightarrow \nabla(\mathcal{P}^C(U)) \quad \text{in } [L^q(\Omega)]^{n \times N}, \quad \text{if } q \in (1, \infty), \quad (\text{C2b})$$

$$\mathcal{L}^n \llcorner \nabla(\mathcal{F}_m^C(U^m)) \xrightarrow{*} \mathcal{L}^n \llcorner \nabla(\mathcal{P}^C(U)) \quad \text{in } [\mathcal{M}(\Omega)]^{n \times N}, \quad \text{if } q = 1. \quad (\text{C2c})$$

In particular, (C2a), (C2b), (C2c) together with (C1) mean that  $\nabla(\mathcal{F}_m^C(U^m)) \rightarrow \nabla(\mathcal{P}^C(U))$  in the weak topology  $\mathcal{T}_{1;p,q}^w$  of  $[W_0^{1;p,q}(\Omega)]^N$  in (Afs). Also, in view of  $(\diamond)$ ,  $|\nabla(\mathcal{F}_m^C(U^m))| = |\nabla U^m|$  for all  $m = 1, 2, \dots$ .

**Remark 4.7.** The conclusions (C2a), (C2b), (C2c) do not contradict  $|\nabla(\mathcal{F}_m^C(U^m))| = |\nabla U^m|$ , since bounded sequences of constant norm may well be weakly zero. A related example is given by the sequence of Foldings of  $u_0 \in W^{1,\infty}(0,2)$  defined inductively by  $u_0(x) := x\chi_{(0,1)}(x) + (2-x)\chi_{(1,2)}(x)$  and  $u_{m+1} := u_m\chi_{\{u_m < 2^{-(m+1)}\}} + (2^{-m} - u_m)\chi_{\{u_m > 2^{-(m+1)}\}}$  in  $W^{1,\infty}(0,2)$ . We have  $\|u_m\|_{W^{1,\infty}(0,2)} \leq 1$ . Since  $\|u_m\|_{L^\infty(0,2)} \leq 2^{-m-1}$ , the weak\* limit is zero. Thus  $u_m \xrightarrow{*} 0$  in  $W^{1,\infty}(0,2)$ , which means  $u_m \rightarrow 0$  in  $L^\infty(0,2)$  and  $(u_m)_x \xrightarrow{*} 0$  in  $L^\infty(0,2)$ , although,  $|(u_m)_x| = 1$   $\mathcal{L}^1 \llcorner (0,2)$ -a.e. .



**A representation formula of the essential range (Essrng) of  $\mu$ -measurable functions.** In order to establish Lemma 4.6 and the main theorems 3.1 and 3.7 we need the following pointwise expression of (Essrng). Let  $\mu \in \mathcal{M}(\Omega)$  be Radon with  $U : \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^N$   $\mu$ -measurable. Let  $\Omega \setminus S_U^\mu$  be the  $\mu$ -approximate continuity set of  $U$  and  $\{U^\mu(x) : x \in \Omega \setminus S_U^\mu\}$  the set of  $\mu$ -approximate limits:

$$\Omega \setminus S_U^\mu := \left\{ x \in \Omega \mid \exists U^\mu(x) \in \mathbb{R}^N : \forall \varepsilon > 0, \lim_{r \rightarrow 0} \frac{\mu(\mathbb{B}_r(x) \cap \{|U - U^\mu(x)| \geq \varepsilon\})}{\mu(\mathbb{B}_r(x))} = 0 \right\}. \quad (\text{AppC}^0)$$

In this case we set

$$U^\mu(x) := \mu - \text{app } \lim_{z \rightarrow x} U(z), \quad \text{for all } x \in \Omega \setminus S_U^\mu. \quad (\text{AppLim})$$

By [[E-G], Th. 3, Sec. 1.7] and trivial modifications,  $\mu(S_U^\mu) = 0$  and  $U^\mu$  is  $\mu$ -a.e. defined, measurable and invariant under modifications of  $U$  on nullsets. If  $U \in [L_{\text{loc}}^\infty(\Omega, \mu)]^N$ , then  $U^\mu : \Omega \setminus S_U^\mu \rightarrow \mathbb{R}^N$  coincides with the Besicovitch  $\mu$ -precise representative of  $U$ . The proof of Ambrosio-Fusco-Palara [[A-F-P], Prop.3.65] for  $\mu = \mathcal{L}^n$  easily adapts to any  $\mu$ . When  $\mu = \mathcal{L}^n$ , we set  $U^{\mathcal{L}^n} = U^*$  and  $S_U^\mu = S_U^*$ .

**Lemma 4.8.** *Let  $\mu \in \mathcal{M}(\Omega)$  be positive. For any  $U : \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^N$  and  $A \subseteq \Omega$  both  $\mu$ -measurable with  $\mu(A) > 0$ , we have the following representation formula for (Essrng):*

$$\mu - \text{ess } U(A) = \overline{U^\mu(\text{int}_\mu(A) \setminus S_U^\mu)}. \quad (\text{Essrng})$$

where the measure theoretic interior  $\text{int}_\mu(A)$  of the set  $A$  with respect to  $\mu$  is defined via the Lebesgue set of  $\chi_A$ :  $\text{int}_\mu(A) := \Omega \setminus S_{\chi_A}^\mu \cap \{(\chi_A)^\mu = 1\}$ . Similarly, the  $\mu$ -exterior is  $\text{ext}_\mu(A) := \Omega \setminus S_{\chi_A}^\mu \cap \{(\chi_A)^\mu = 0\}$ .

Lemma 4.8 implies that when  $U \in [W_{\text{loc}}^{1;p,q}(\mathbb{R}^n)]^N$  and  $K \subseteq \overline{\Omega}$  is bounded with  $\mathcal{H}^{n-1}(K) > 0$ , we can represent the convex hull (CO) by  $\text{app } \overline{\text{co}}(U(K)) = \overline{\text{co}}(U^*(K \setminus S_U^*))$ . The same holds if  $U \in [W^{1;p,q}(\Omega)]^N$  and  $\partial\Omega$  is Lipschitz. Also, an equivalent definition of (EssCo) is  $\mu - \text{ess } \overline{\text{co}}(U(A)) = \overline{\text{co}}(\mu - \text{ess } U(A))$ .

**Remark 4.9.** *When dealing with compositions of nonlinear continuous with measurable functions, as we repeatedly do in this paper, we will consider them as defined only on the possibly smaller initial approximate continuity set. In particular, when  $U \in [L_{\text{loc}}^1(\Omega)]^N$  and  $F \in [W^{1,\infty}(\mathbb{R}^N)]^N$ , the estimate*

$$\int_{\mathbb{B}_r(x)} |F(U) - F(U^*(x))| d\mathcal{L}^n \leq \text{Lip}[F, \mathbb{R}^N] \int_{\mathbb{B}_r(x)} |U - U^*(x)| d\mathcal{L}^n$$

implies  $\Omega \setminus S_U^* \subseteq \Omega \setminus S_{F(U)}^*$  and  $(F(U))^* = F(U^*)$  there. Thus, we may define nonlinear compositions by  $F(U) := (\chi_{\Omega \setminus S_U^*})^{-1} F(U^*)$ , by which we mean we set  $F(U) := \infty$  on the (harmless) nullset  $S_U^*$ . This practice saves a lot of labor when considering pointwise properties, since we avoid considering  $(F(U))^*$ .

## 5 Proofs.

**Proof of Lemma 4.1.** (i) We consider the subset of the sphere  $C^\sharp := \{\xi \in \{\hat{u} : u \in \partial C\} \subseteq \mathbb{S}^{N-1}$ . Generally,  $\partial C$  is a  $W^{1,\infty}$  hypersurface and  $u \mapsto \{\lambda \hat{u} : \lambda > 0\}$  is the outwards normal cone, which is 1-dimensional  $\mathcal{H}^{N-1} \llcorner \partial C$ -a.e.. The stereographic projection  $\pi_N : \mathbb{S}^{N-1} \setminus \{e_N\} \rightarrow \mathbb{R}^{N-1}$  given by  $(\xi_1, \dots, \xi_N) \mapsto (\frac{\xi_1}{\xi_N-1}, \dots, \frac{\xi_{N-1}}{\xi_N-1})$  is a diffeomorphism. Let  $R : \mathbb{S}^{N-1} \rightarrow \text{O}(N, \mathbb{R})$  be the reflection map  $\xi \mapsto R_\xi$ . Then, the mapping  $R \circ \pi_N^{-1} : \mathbb{R}^{N-1} \rightarrow \text{O}(N, \mathbb{R})$  is  $C^1$ , since for any  $\varepsilon > 0$ ,

$$\sup_{|z|, |u| \leq 1} \left| \frac{1}{\varepsilon} (R_{\xi+\varepsilon z} u - R_\xi u) + 2[(u \cdot \xi)z + (u \cdot z)\xi] \right| \leq 2\varepsilon.$$

Restricting  $R \circ \pi_N^{-1}$  on  $\pi_N(C^\sharp \setminus \{e_N\}) \subseteq \mathbb{R}^{N-1}$ , we obtain

$$\mathcal{H}^{N-1} \left( \mathcal{R}(C) \setminus (R \circ \pi_N^{-1}(\pi_N(C^\sharp \setminus \{e_N\}))) \right) = 0.$$

(ii) For any  $u \in \partial C$ ,  $\{\hat{u}\}$  is geodesically convex on  $\mathbb{S}^{N-1}$ , since for any  $\xi, \eta \in \{\hat{u}\}$ , there is an arc  $\phi : [0, 1] \rightarrow \{\hat{u}\}$  with  $\phi(0) = \xi, \phi(1) = \eta$  given by

$$\lambda \mapsto \phi(\lambda) := \frac{\lambda\eta + (1-\lambda)\xi}{|\lambda\eta + (1-\lambda)\xi|}.$$

$\phi$  is well defined since not both  $\xi$  and  $-\xi$  belong to  $\{\hat{u}\}$ , otherwise for all  $v \in C$ ,  $\xi \cdot (u - v) = 0$  and then  $C \subseteq u + (\text{span}[\xi])^\perp$ , contradicting  $\text{int}(C) \neq \emptyset$ . We obtain that  $C^\#$  is path connected, since  $C^\# = \bigcup\{\{\hat{u}\} : u \in \partial C\}$  and  $\text{dist}(\{\hat{u}\}, \{\hat{v}\}) \rightarrow 0$  when  $|u - v| \rightarrow 0$ . By continuity of  $R$ ,  $\mathcal{R}(C) \setminus \{I\}$  is path connected set of reflections in  $O(N, \mathbb{R}) \cap \{det = -1\}$ .

For (iii) and (iv), we first show that the set  $\mathcal{R}(\mathbb{B}^N) \setminus \{I\}$  is diffeomorphic to  $\mathbb{RP}^{N-1}$ . The latter manifold consists of equivalence classes  $\{[v]_\sim : v \in \mathbb{S}^{N-1}\}$  where  $v' \sim v'' \Leftrightarrow v'' = -v'$ . That is, the antipodal points of  $\mathbb{S}^{N-1}$  are identified.  $R$  induces a mapping from the quotient  $\mathbb{RP}^{N-1} = \mathbb{S}^{N-1}/\sim$  to  $\mathcal{R}(\mathbb{B}^N) \setminus \{I\}$  given by  $[v]_\sim \mapsto R_{\hat{v}}$ . It is well defined on equivalence classes, bijective and  $v'' \sim v' \Leftrightarrow v'' = -v' \Leftrightarrow R_{\hat{v}'} = R_{-\hat{v}'}$ . Standard geometric arguments imply that it is a diffeomorphism w.r.t. the natural smooth structures induced by quotient and inclusion. We now show that for any  $C \subseteq \mathbb{R}^N$  compact convex,  $\mathcal{R}(C) \setminus \{I\} = \mathcal{R}(\mathbb{B}^N) \setminus \{I\}$  and for merely closed  $C$  we have the injection  $\mathcal{R}(C) \hookrightarrow \mathcal{R}(\mathbb{B}^N)$ . Since  $\mathcal{R}(C)$  is invariant under translations and dilations of  $C$ , we may assume that  $\mathbb{B}^N \subset\subset C$ ,  $\mathbb{B}^N = \mathbb{B}_1^N(0)$ . Let  $H = u + H_{\hat{u}} \in \mathcal{S}(C)$ . Then,  $H' := (\hat{u} - u) + H_{\hat{u}} \in \mathcal{S}(\mathbb{B}^N)$  and as such,  $\mathcal{R}(C) \subseteq \mathcal{R}(\mathbb{B}^N)$ .

For the reverse inclusion we employ an argument that does not require convexity and applies to compact connected Lipschitz domains. Let  $\xi \in \partial \mathbb{B}^N = \mathbb{S}^{N-1}$  with  $H = \xi + (\text{span}[\xi])^\perp$  in  $\mathcal{S}(\mathbb{B}^N)$ . For  $t \in \mathbb{R}$ , set  $H_t := t\xi + (\text{span}[\xi])^\perp$  and  $h(t) := \mathcal{H}^{N-1}(H_t \cap C)$ .

**Claim 5.1.**  $h \in C_c^0(\mathbb{R})$  and there exist  $t_{\min} < 0, t_{\max} > 0$  such that  $\text{supp}(h) = [t_{\min}, t_{\max}]$ .

**Proof of Claim 5.1.** The translates  $H(t) := (H_t \cap C) - t\xi$  of the slices  $H_t \cap C$  satisfy  $H(t) \subseteq (\text{span}[\xi])^\perp \cong \mathbb{R}^{N-1}$  and also  $\mathcal{H}^{N-1}(H_t \cap C) = \mathcal{H}^{N-1}(H(t))$ . If  $t \rightarrow t_0$ , then  $H(t) \rightarrow H(t_0)$  in  $\mathcal{H}^{N-1} \llcorner (\text{span}[\xi])^\perp$  measure, i.e.  $\chi_{H(t)} \rightarrow \chi_{H(t_0)}$  in  $L^1((\text{span}[\xi])^\perp, \mathcal{H}^{N-1})$ . Thus, we obtain  $h(t) \rightarrow h(t_0)$  as  $t \rightarrow t_0$  and  $h(t) = 0$  for  $t > \text{diam}(C)$ . Let  $[t_{\min}, t_{\max}] := \overline{\text{co}}(\text{supp}(h))$ . By path connectivity and compactness of  $C$ ,  $t_{\min}\xi, t_{\max}\xi \in \partial C$  and there exists a path  $\psi : [t_{\min}, t_{\max}] \rightarrow C$  with  $\psi(t_{\min}) = t_{\min}\xi$  and  $\psi(t_{\max}) = t_{\max}\xi$ . Since  $\text{int}(C) \neq \emptyset$ , for all  $t \in (t_{\min}, t_{\max})$  there is a ball  $\mathbb{B}_{\varepsilon_t}^N(\psi(t)) \subseteq \text{int}(C)$ . Thus,  $h(t) = \mathcal{H}^{N-1}(H_t \cap C) \geq \mathcal{H}^{N-1}(H_t \cap \mathbb{B}_{\varepsilon_t}^N(\psi(t))) > 0$  and the Claim 5.1 follows.  $\square$

The next Claim completes the proof of Lemma 4.1.

**Claim 5.2.** The affine hyperplanes  $H_{t_{\max}}$  and  $H_{t_{\min}}$  belong to  $\mathcal{S}(C)$ . Thus, we obtain  $\mathcal{R}(\mathbb{B}^N) \subseteq \mathcal{R}(C)$ .

**Proof of Claim 5.2.** Every  $u \in C$  can be written as  $u = t\xi + z$ , with  $z \in (\text{span}[\xi])^\perp$  and  $t \in [t_{\min}, t_{\max}]$ . Thus,  $\xi \cdot (t_{\max}\xi - u) = t_{\max}|\xi|^2 - \xi \cdot (t\xi + z) = (t_{\max} - t)|\xi|^2 - \xi \cdot z = t_{\max} - t \geq 0$ . The other is similar.  $\square$

Note that closeness of  $\mathcal{R}(C)$  may fail if  $\text{diam}(C) = \infty$ : suffices to take  $C := \{(u, v) \in \mathbb{R}^2 : uv \geq 1 \text{ if } 0 < u \leq 1 \text{ and } (2-u)v \geq 1 \text{ if } 1 \leq u < 2\}$ . Then, as  $u \rightarrow 0$   $(u, u^{-1}) \rightarrow \infty$  but  $(u, u^{-1}) \rightarrow (-1, 0) \notin \mathcal{R}(C)$ .

**Proof of Lemma 4.4.** Since  $C$  is fixed, we drop superscripts. We shall first prove the existence of  $\mathcal{F}_\lambda$  for obtuse simplices satisfying (O) and then deduce the case of general obtuse convex sets by approximation.



For, assume  $C = \bigcap \{H_j^+ : 1 \leq j \leq K\}$  with  $K \geq N + 1$  and  $H_j = u_j + (\text{span}[\xi_j])^\perp$  with (O) being satisfied by the sides  $H_j^*$ . If  $\lambda > 1$ ,  $C \subseteq \lambda C$  and  $\lambda C = \bigcap \{\lambda H_j : 1 \leq j \leq K\}$  since  $\lambda H_j = \lambda u_j + (\text{span}[\xi_j])^\perp$ . We choose  $\xi_{\min}$  on  $\partial C = \bigcap \{H_j : 1 \leq j \leq K\}$  and  $\xi_{\max}$  on  $\bigcup \{H_j : 1 \leq j \leq K\}$  such that


$$|\xi_{\min}| = \min_{1 \leq j \leq K} \left( \min_{u \in H_j} |u| \right), \quad |\xi_{\max}| = \max_{1 \leq j \leq K} \left( \min_{u \in H_j} |u| \right). \quad (\text{minmax})$$

The points  $\xi_{\min}$  and  $\xi_{\max}$  always exist and are normal to  $\bigcup \{H_j : 1 \leq j \leq K\}$ , but possibly are non unique or  $|\xi_{\min}| = |\xi_{\max}|$ . By (minmax),  $\mathbb{B}_{|\xi_{\min}|}(0) \subseteq C$  and this is the largest such ball. We set:

$$T := \frac{1}{|\xi_{\min}|} \max \left\{ a, \sup_{u \in C} |u| \right\} = \frac{1}{|\xi_{\min}|} \sup_{u \in A \cup C} |u|.$$

The dilation  $u \mapsto Tu$  maps  $C$  onto  $TC$  and, by the choice of  $T$ ,

$$A \subseteq A \cup C \subseteq \mathbb{B}_{T|\xi_{\min}|}(0) \subseteq TC.$$



We fix two numbers  $t, \tau > 0$  with  $1 < t < \tau \leq T$ , chosen such that

$$t = \frac{\tau |\xi_{\max}| + |\xi_{\min}|}{|\xi_{\min}| + |\xi_{\max}|}. \quad (**)$$

By (minmax), the min distance between the sides of  $\partial C$  and  $\partial(tC)$  is realized in the  $\xi_{\min}$  direction, while the max distance between the sides of  $\partial(\tau C)$  and  $\partial(tC)$  in the  $\xi_{\max}$  direction. Thus, (\*\*) rearranged says

$$\min_{1 \leq j \leq K} \left( \min_{u \in H_j} |tu - u| \right) = \max_{1 \leq j \leq K} \left( \min_{u \in H_j} |\tau u - tu| \right).$$

We take  $s \in [1, T]$  and  $j \in \{1, \dots, K\}$ . The affine reflection  $R_j^s \in \text{AO}(N, \mathbb{R})$  with respect to  $sH_j$  is given by  $R_j^s(u) = R_{\xi_j} u + 2s(u_j \cdot \xi_j)\xi_j$ ,  $R_{\xi_j} \in \mathcal{R}^*(C)$ . For any  $s \in [1, T]$  and  $j \in \{1, \dots, K\}$ , we define

$$\mathcal{F}_j^s(u) := \begin{cases} u, & u \in sH_j^+ \\ R_j^s(u), & u \notin sH_j^+ \end{cases}, \quad \mathcal{F}_j^s : \mathbb{R}^N \longrightarrow \mathbb{R}^N, \quad (\mathcal{F}I)$$

and


$$\mathcal{F}^s := \mathcal{F}_K^s \circ \dots \circ \mathcal{F}_1^s, \quad \mathcal{F}^s : \mathbb{R}^N \longrightarrow \mathbb{R}^N. \quad (\mathcal{F}II)$$

Each  $\mathcal{F}_j^s$  is non-expansive in  $[W_{\text{loc}}^{1,\infty}(\mathbb{R}^N)]^N$  with  $\mathcal{F}_j^s|_{sC} = Id_{sC}$  and  $\text{Lip}[\mathcal{F}_j^s] = 1$ . Thus, we have that  $\mathcal{F}^s|_{sC} = Id_{sC}$  and  $\text{Lip}[\mathcal{F}^s] = 1$  being in  $[W_{\text{loc}}^{1,\infty}(\mathbb{R}^N)]^N$ .

**Claim 5.3.** Assume that  $t, \tau$  are in  $[1, T]$  and they satisfy (\*\*). Then, for all  $j = 1, \dots, K$ , we have

$$\mathcal{F}_j^t(\tau C \setminus \text{int}(tH_j^+)) \subseteq \tau C \cap (tH_j^+ \setminus \text{int}(H_j^+)).$$

**Proof of Claim 5.3.** For any  $B \subseteq \tau C$ , (F1) implies that  $\mathcal{F}_j^t(B) = (B \cap tH_j^+) \cup R_j^t(B \setminus tH_j^+)$ . First note that by (\*\*),  $\text{dist}(\tau H_j^+, tH_j^+) \leq \text{dist}(tH_j^+, H_j^+)$ ,  $j = 1, \dots, K$ , thus  $\mathcal{F}_j^t$  maps the zone  $\tau H_j^+ \setminus \text{int}(tH_j^+)$  into the zone  $tH_j^+ \setminus \text{int}(H_j^+)$ . We may first normalize up to an element of  $\text{AO}(N, \mathbb{R})$  to  $\xi_j = e_N$ ,  $tH_j = \{u_N = 0\}$ .



Let  $u = (u', u_N) \in \tau C \setminus \text{int}(tH_j^+)$ . Upon identifying the affine reflections with their standard matrix, we have  $R_j^t(u) = (u', -u_N)$  and  $R_j^t(u)$  belongs to  $tH_j^+ \setminus \text{int}(H_j^+) = \{(1-t)|u_j| \leq u_N \leq 0\}$ . By (O), it suffices to show that  $R_j^t(u)$  is into  $\cap \{\tau H_i^+ : i \neq j, H_i^* \cap H_j^* \neq \emptyset\}$ , for all the neighboring sides  $H_i^*$  to  $H_j^*$ . Fix such an  $H_i^*$ . Then,  $\tau H_i = (\text{span}[\xi_i])^\perp + \tau u_i$  and by a further translation we assume  $u_i \equiv 0$  (of course, the rest defining points  $\{u_l : l \neq i\}$  of  $C$  necessarily change, but this affects nothing). We obtain that  $\tau H_i = \{z \in \mathbb{R}^N : z \cdot \xi_i = 0\}$  and  $\tau H_i^+$  is the subgraph of the linear functional  $z \mapsto z \cdot \xi_i$ . Since  $u \in \tau C = \cap \{\tau H_j^+ : 1 \leq j \leq K\} \subseteq \tau H_i^+$ , we have  $u \cdot \xi_i \leq 0$  and this gives

$$\begin{aligned} u' \cdot \xi_i' &= -(u \cdot e_N)(\xi_i \cdot e_N) + u \cdot \xi_i \\ &\leq -(u \cdot e_N)(\xi_i \cdot e_N). \end{aligned}$$

Thus, utilizing that  $u \in \{u_N \geq 0\}$  and that (O) is equivalent to  $\xi_i \cdot e_N \in [0, 1]$  for all such  $i \neq j$ , we obtain

$$\begin{aligned} (R_j^t(u)) \cdot \xi_i &= u' \cdot \xi_i' + (-u \cdot e_N)(\xi_i \cdot e_N) \\ &\leq -2(u \cdot e_N)(\xi_i \cdot e_N) \leq 0. \end{aligned}$$

Consequently, if  $u \in \tau C \setminus \text{int}(tH_j^+)$ , we have  $\mathcal{F}_j^t(u) \in \tau H_i^+$  for all  $i \neq j$  for which  $H_i^* \cap H_j^* \neq \emptyset$ .  $\square$

**Claim 5.4.** Assume that  $t, \tau$  belong to  $[1, T]$  and they satisfy (\*\*). Then, we have

$$\begin{cases} (i) & \mathcal{F}^t(\tau C \setminus \text{int}(tC)) \subseteq tC \setminus \text{int}(C), \\ (ii) & \mathcal{F}^t(\tau C) \subseteq tC. \end{cases}$$

**Proof of Claim 5.4.** (ii) is a consequence of (i) since  $\mathcal{F}^t|_{tC} = \text{Id}_{tC}$ , so it suffices to prove (i). Let  $u \in \tau C \setminus \text{int}(tC)$  be a fixed point. We first show that  $\mathcal{F}^t(u) \notin \text{int}(C)$ . Let  $k \in \{1, \dots, K\}$  be the first in order index for which  $u \in \tau C \setminus \text{int}(tH_k^+)$ . By (F1), all  $\mathcal{F}_1^t, \dots, \mathcal{F}_{k-1}^t$  leave  $u$  invariant, hence  $\mathcal{F}^t(u) = \mathcal{F}_K^t(\dots \mathcal{F}_k^t(u))$ . By Claim 5.3,  $\mathcal{F}_k^t(u) \in \tau C \cap (tH_k^+ \setminus \text{int}(H_k^+))$  and in particular  $\mathcal{F}^t(u) \notin \text{int}(C)$ . Let  $l \geq k+1$  be the next in order index for which

$$\mathcal{F}_k^t(u) \in \tau C \cap (tH_k^+ \setminus \text{int}(H_k^+)) \setminus \text{int}(tH_l^+) \subseteq \tau C \setminus \text{int}(tH_l^+).$$

For all the intermediate indices  $k+1 \leq i < l$ ,  $\mathcal{F}_i^t$  leaves  $\mathcal{F}_k^t(u)$  invariant and

$$\mathcal{F}_l^t(\mathcal{F}_{l-1}^t \dots \mathcal{F}_i^t \dots (\mathcal{F}_k^t(u))) = \mathcal{F}_l^t(\mathcal{F}_k^t(u)).$$

By Claim 5.3,  $\mathcal{F}_l^t(\mathcal{F}_k^t(u)) \in \tau C \cap (tH_l^+ \setminus \text{int}(H_l^+))$ , thus  $\mathcal{F}_l^t(\mathcal{F}_k^t(u)) \notin \text{int}(C)$ . Continuing till  $K$ , in view on (FII) we obtain  $\mathcal{F}^t(u) \notin \text{int}(C)$ . We now show that  $\mathcal{F}^t(u) \in \text{int}(tC)$ . Let again  $k \in \{1, \dots, K\}$  be the first in order index for which  $u \in \tau C \setminus tH_k^+$ . Then,  $\mathcal{F}^t(u) = \mathcal{F}_K^t(\dots \mathcal{F}_k^t(u))$ . Let  $H_l^*$ ,  $l \geq k+1$ , be the first in order neighboring side of  $H_k^*$ . By Claim 5.3,  $\mathcal{F}_k^t(u) \in \tau C \cap (tH_k^+ \setminus \text{int}(H_k^+))$ . If  $\mathcal{F}_k^t(u) \in tH_l^+ \cap \tau C$ , we proceed to the next neighboring side. Otherwise, if  $\mathcal{F}_k^t(u) \in \tau C \setminus \text{int}(tH_l^+)$ , the exact same argument in the proof of Claim 5.3 with  $tH_k$  in the place of  $tH_j$  and  $tH_l$  in the place of  $\tau H_i$  shows that

$$\mathcal{F}_l^t(\mathcal{F}_k^t(u)) \in \tau C \cap (tH_k^+ \setminus \text{int}(H_k^+)) \cap (tH_l^+ \setminus \text{int}(H_l^+)).$$

In view of (FII), continuing till  $K$  we obtain  $\mathcal{F}^t(u) \in tC \cap \{tH_j^+ \setminus \text{int}(H_j^+) : 1 \leq j \leq K\} \subseteq tC$  as desired.  $\square$

Now we iterate Claim 5.4. We define inductively:

$$\begin{cases} t_0 := T \\ t_{k+1} := \frac{t_k |\xi_{\max}| + |\xi_{\min}|}{|\xi_{\min}| + |\xi_{\max}|}. \end{cases} \quad (***)$$

**Claim 5.5.** *The sequence  $(t_k)_{k=1}^\infty$  defined by (\*\*\*) is strictly decreasing to  $1^+$  as  $k \rightarrow \infty$ , and*

$$t_k = \left( \frac{|\xi_{\max}|}{|\xi_{\max}| + |\xi_{\min}|} \right)^k T + \left( \frac{|\xi_{\min}|}{|\xi_{\max}| + |\xi_{\min}|} \right) \sum_{j=0}^{k-1} \left( \frac{|\xi_{\max}|}{|\xi_{\max}| + |\xi_{\min}|} \right)^j, \quad k = 1, 2, \dots$$

**Proof of Claim 5.5.** Follows by induction on (\*\*\*) and the geometric series.  $\square$

Given  $\lambda > 1$ , we choose  $k^* \in \mathbb{N}$ :  $t_{k^*} < \lambda$ . Then,  $t_{k^*}C \subseteq \lambda C$ . We define:

$$\mathcal{F}_\lambda^C := \mathcal{F}^{t_{k^*}} \circ \dots \circ \mathcal{F}^{t_1}. \quad (\text{FIII})$$

The following Claim completes the proof of the Lemma 4.4.

**Claim 5.6.** *The Folding Map  $\mathcal{F}_\lambda^C$  given by (FI), (FII), (FIII) is non-expansive locally Lipschitz in  $[W_{loc}^{1,\infty}(\mathbb{R}^N)]^N$  with  $\text{Lip}[\mathcal{F}_\lambda^C] = 1$ , it is composed by elements of  $\mathcal{R}^*(C)$  and satisfies (\*).*

**Proof of Claim 5.6.** By Claims 5.4, 5.5, for all  $k = 1, 2, \dots$  we have

$$\begin{cases} \mathcal{F}^{t_k}(t_{k-1}C \setminus \text{int}(t_k C)) \subseteq t_k C \setminus \text{int}(C) \\ \mathcal{F}^{t_k}(t_{k-1}C) \subseteq t_k C. \end{cases}$$

It suffices to verify (\*), since the rest follow directly by (FI), (FII), (FIII). Since  $A \subseteq TC = t_0 C$ ,

$$\begin{aligned} \mathcal{F}_\lambda^C(A) &\subseteq \mathcal{F}^{t_{k^*}}(\dots \mathcal{F}^{t_1}(t_0 C)) \subseteq \mathcal{F}^{t_{k^*}}(\dots \mathcal{F}^{t_2}(t_1 C)) \\ &\subseteq \mathcal{F}^{t_{k^*}}(\dots \mathcal{F}^{t_3}(t_2 C)) \subseteq \dots \subseteq t_{k^*} C \\ &\subseteq \lambda C. \end{aligned}$$

Now, since  $A \setminus C \subseteq A \setminus \text{int}(C) \subseteq TC \setminus \text{int}(C) = t_0C \setminus \text{int}(C)$ , utilizing (FI), (FII), (FIH) we have

$$\begin{aligned} \mathcal{F}^{t_1}(A \setminus C) &\subseteq \mathcal{F}^{t_1}(t_0C \setminus \text{int}(C)) \\ &= \mathcal{F}^{t_1}\left([(t_0C \setminus \text{int}(C)) \cap \text{int}(t_1C)] \cup [(t_0C \setminus \text{int}(C)) \setminus \text{int}(t_1C)]\right) \\ &\subseteq (t_1C \setminus \text{int}(C)) \cup \mathcal{F}^{t_1}(t_0C \setminus \text{int}(t_1C)) \\ &\subseteq (t_1C \setminus \text{int}(C)) \cup (t_1C \setminus \text{int}(C)) \\ &= t_1C \setminus \text{int}(C), \end{aligned}$$

$$\begin{aligned} \mathcal{F}^{t_2}(\mathcal{F}^{t_1}(A \setminus C)) &\subseteq \mathcal{F}^{t_2}(t_1C \setminus \text{int}(C)) \\ &= \mathcal{F}^{t_2}\left([(t_1C \setminus \text{int}(C)) \cap \text{int}(t_2C)] \cup [(t_1C \setminus \text{int}(C)) \setminus \text{int}(t_2C)]\right) \\ &\subseteq (t_2C \setminus \text{int}(C)) \cup \mathcal{F}^{t_2}(t_1C \setminus \text{int}(t_2C)) \\ &\subseteq (t_2C \setminus \text{int}(C)) \cup (t_2C \setminus \text{int}(C)) \\ &= t_2C \setminus \text{int}(C), \end{aligned}$$

$\vdots$

$$\begin{aligned} \mathcal{F}_\lambda^C(A \setminus C) &\subseteq \mathcal{F}^{t_k^*}(\dots \mathcal{F}^{t_1}(A \setminus \text{int}(C))) \subseteq \dots \subseteq t_k^*C \setminus \text{int}(C) \\ &\subseteq \lambda C \setminus \text{int}(C). \end{aligned}$$

The Claim 5.6 has been established.  $\square$

The proof of Lemma 4.4 follows by approximating general obtuse convex sets by obtuse simplices.  $\square$

**Proof of Lemma 4.5.** We temporarily assume that  $C$  is obtuse satisfying (O1). We first show that there exists a non-expansive Lipschitz function  $\mathcal{F}^C : \mathbb{R}^N \rightarrow C$  in  $[W^{1,\infty}(\mathbb{R}^N)]^N$  with  $\text{Lip}[\mathcal{F}^C] = 1$  and

$$\begin{cases} \mathcal{F}^C(E \cap C) = E \cap C, & E \subseteq \mathbb{R}^N \\ \mathcal{F}^C(\mathbb{R}^N \setminus C) \subseteq \partial C, \end{cases} \quad (\diamond)$$

which is a weak\* sequential limit of  $\mathcal{F}_\lambda^C$  as  $\lambda \rightarrow 1^+$  in  $[W_{\text{loc}}^{1,\infty}(\mathbb{R}^N)]^N$ . Set  $B_m := \mathbb{B}_m(0)$ ,  $m = 1, 2, \dots$ . By Lemma 4.4, for any  $\lambda > 1$ , there is a Folding Map  $\mathcal{F}_\lambda^{C,m} : \mathbb{R}^N \rightarrow \mathbb{R}^N$ , with  $\text{Lip}[\mathcal{F}_\lambda^{C,m}] = 1$  in  $[W_{\text{loc}}^{1,\infty}(\mathbb{R}^N)]^N$  satisfying

$$\begin{cases} \mathcal{F}_\lambda^{C,m}(E \cap C) = E \cap C, & E \subseteq \mathbb{R}^N \\ \mathcal{F}_\lambda^{C,m}(B_m \setminus C) \subseteq \lambda C \setminus \text{int}(C). \end{cases}$$

Note that  $\mathcal{F}_\lambda^{C,m}(B_m) \subseteq \lambda C$  for all  $m$ . We take  $\lambda_m := 1 + \frac{1}{m}$  and fix  $k \in \mathbb{N}$ . Then, for  $m \geq k$ ,

$$\mathcal{F}_{1+\frac{1}{m}}^{C,m}(B_k) \subseteq \left(1 + \frac{1}{m}\right)C, \quad (\text{i})$$

$$\text{ess sup}_{B_k} |\nabla \mathcal{F}_{1+\frac{1}{m}}^{C,m}| \leq \text{Lip}[\mathcal{F}_{1+\frac{1}{m}}^{C,m}] = 1, \quad (\text{ii})$$

$$\mathcal{F}_{1+\frac{1}{m}}^{C,m}(E \cap C) = E \cap C, \quad E \subseteq \mathbb{R}^N, \quad (\text{iii})$$

$$\mathcal{F}_{1+\frac{1}{m}}^{C,m}(B_k \setminus C) \subseteq \left(1 + \frac{1}{m}\right)C \setminus \text{int}(C). \quad (\text{iv})$$

Thus, by (i) and (ii), we have

$$\left\| \mathcal{F}_{1+\frac{1}{m}}^{C,m} \right\|_{[W^{1,\infty}(B_k)]^N} \leq 2 \max_{u \in C} |u| + 1 \quad \text{for all } m \geq k,$$

and by weak\* compactness of the local Frechét space  $[W_{\text{loc}}^{1,\infty}(\mathbb{R}^N)]^N$ , we can extract a (renamed) subsequence such that  $\mathcal{F}_{1+\frac{1}{m}}^{C,m} \xrightarrow{*} \mathcal{F}^C$  as  $m \rightarrow \infty$ , for some  $\mathcal{F}^C \in [W_{\text{loc}}^{1,\infty}(\mathbb{R}^N)]^N$ . By standard compactness reasoning,  $\mathcal{F}_{1+\frac{1}{m}}^{C,m} \rightarrow \mathcal{F}^C$  in  $[L_{\text{loc}}^\infty(\mathbb{R}^N)]^N$ . Passing to the limit to (iii) and (iv), we obtain

$$\mathcal{F}^C(E \cap C) = E \cap C \quad \text{and} \quad \mathcal{F}^C(B_k \setminus C) \subseteq \partial C.$$

Hence,  $\mathcal{F}^C(B_k) \subseteq C$ . Since  $\mathbb{R}^N = \bigcup_{k \geq 1} B_k$ , we get  $\mathcal{F}^C(\mathbb{R}^N) = \bigcup_{k \geq 1} \mathcal{F}^C(B_k) \subseteq C$  and  $\mathcal{F}^C(\mathbb{R}^N \setminus C) = \bigcup_{k \geq 1} \mathcal{F}^C(B_k \setminus C) \subseteq \partial C$ . Thus, we obtain that  $\mathcal{F}^C \in [L^\infty(\mathbb{R}^N)]^N$  and  $(\diamond)$  follows in this case.

Now we drop obtuseness (O1). Let  $(C_m)_1^\infty \subseteq \mathbb{R}^N$  be a decreasing sequence of smooth compact convex sets which approximate  $C$  from the outside:  $C_m \supseteq C_{m+1} \supseteq \dots \supseteq C$ . For example, we may take  $C_m := \{\text{dist}(\cdot, C) \leq \frac{1}{m}\}$ . For each  $m \in \mathbb{N}$ , there is an  $\mathcal{F}^{C_m} : \mathbb{R}^N \rightarrow \mathbb{R}^N$  with  $\mathcal{F}^{C_m}(E \cap C_m) = E \cap C_m$ , for any  $E \subseteq \mathbb{R}^N$  and  $\mathcal{F}^{C_m}(\mathbb{R}^N \setminus C_m) \subseteq \partial C_m$ . For fixed  $k \in \mathbb{N}$  and  $m \geq k$ , we have

$$\begin{aligned} \mathcal{F}^{C_m}(B_k) &\subseteq \mathcal{F}^{C_m}(B_k \cap C_m) \cup \mathcal{F}^{C_m}(B_k \setminus C_m) \subseteq C_m \cup \partial C_m \subseteq C_1, \\ \text{ess sup}_{B_k} |\nabla \mathcal{F}^{C_m}| &\leq \text{Lip}[\mathcal{F}^{C_m}] = 1. \end{aligned}$$

The weak\* compactness of  $[W_{\text{loc}}^{1,\infty}(\mathbb{R}^N)]^N$ , implies again that up to a subsequence  $\mathcal{F}^{C_m} \xrightarrow{*} \mathcal{F}^C$  as  $m \rightarrow \infty$  and we again obtain  $(\diamond)$ . Finally, we recall from Lemma 4.1 (iii) that the sets  $\mathcal{R}(C)$  and  $\mathcal{R}(C_m)$  of  $C$  and the approximating sequence  $C_m$  respectively satisfy  $\mathcal{R}(C) \setminus \{I\} = \mathcal{R}(C_m) \setminus \{I\} \cong \mathbb{R}^{N-1}$  for all  $m \in \mathbb{N}$ .

We now show that  $\mathcal{F}^C = \mathcal{P}^C$ . Let  $u_0 \in \mathbb{R}^N \setminus C$  be fixed and temporarily assume that  $C$  is smooth. The affine hyperplane  $H_0 := \mathcal{P}^C(u_0) + (\text{span}[u_0 - \mathcal{P}^C(u_0)])^\perp$  belongs to  $\mathcal{S}(C)$ . For all approximating sequences of obtuse simplices  $(C_m)_1^\infty$  satisfying (O), a portion of  $H_0$  near  $\mathcal{P}^C(u_0) \in \partial C$  is always a side of all  $C_m$  for large  $m \in \mathbb{N}$ , as a result of (O1). Recall that  $\mathcal{F}^C(u_0) = \lim_{\lambda \rightarrow 1^+} \mathcal{F}_\lambda^C(u_0)$ , at least sequentially. By (F1), (F2) and (F3) for all  $\lambda > 1$ , we have  $\mathcal{F}_\lambda^C(u_0) \in \lambda C \cap (\mathcal{P}^C(u_0) + \text{span}[u_0 - \mathcal{P}^C(u_0)])$ . Letting  $\lambda \rightarrow 1^+$ , we get  $\mathcal{P}^C(u_0) = \mathcal{F}^C(u_0)$  for any  $u_0 \in \mathbb{R}^N \setminus C$ . Trivially,  $\mathcal{P}^C|_C = \mathcal{F}^C|_C = \text{Id}_C$  as well.

For the general case, approximate  $C$  from the outside by a sequence of smooth compact convex sets  $C_m \supseteq C_{m+1} \supseteq \dots \supseteq C$ . For all  $m \in \mathbb{N}$ ,  $\mathcal{P}^{C_m} = \mathcal{F}^{C_m}$ . Thus, for any  $u_0 \in \mathbb{R}^N$ , by standard properties of  $\mathcal{P}^C$ ,

$$\mathcal{F}^C(u_0) = \lim_{m \rightarrow \infty} \mathcal{F}^{C_m}(u_0) = \lim_{m \rightarrow \infty} \mathcal{P}^{C_m}(u_0) = \mathcal{P}^C(u_0).$$

We complete the proof utilizing Lemma 4.1(iii) to get  $\mathcal{R}(C) = \mathcal{R}(C_m)$  for any sequence  $(C_m)_1^\infty$ .  $\square$

We now establish the representation formula (EssRng).

**Proof of Lemma 4.8.** Let  $u_0 \in \mu - \text{ess } U(A)$ . Then, since  $\mu(S_U^\mu) = \mu(S_{\chi_A}^\mu) = 0$  and  $U = U^\mu$   $\mu$ -a.e., for all  $\varepsilon > 0$  we have

$$\begin{aligned} 0 < \mu\left(A \cap \{|U - u| < \varepsilon\}\right) &= \mu\left(\{|U^\mu - u_0| < \varepsilon\} \cap \text{int}_\mu(A) \setminus S_U^\mu\right) \\ &\leq \mu\left(\left\{\inf\{|U^\mu(x) - u_0| : x \in \text{int}_\mu(A) \setminus S_U^\mu\} < \varepsilon\right\}\right). \end{aligned}$$

Thus,  $u_0 \in \overline{U^\mu(\text{int}_\mu(A) \setminus S_U^\mu)}$  since  $\inf \{|U^\mu(x) - u_0| : x \in \text{int}_\mu(A) \setminus S_U^\mu\} = 0$ . Conversely, let  $u_0$  be in  $\overline{U^\mu(\text{int}_\mu(A) \setminus S_U^\mu)}$ . Then, there is an  $x_0 \in \text{int}_\mu(A) \setminus S_U^\mu$  such that  $|u_0 - U^\mu(x_0)| < \varepsilon$ . Also, for any  $\varepsilon > 0$  small, (AppC<sup>0</sup>) gives that there is a  $\delta > 0$ , such that for all  $r < \delta$ ,

$$\mu(\mathbb{B}_r(x_0) \cap A \cap \{|U - U^\mu(x_0)| < \varepsilon\}) \geq (1 - \varepsilon) \mu(\mathbb{B}_r(x_0) \cap A).$$

Similarly, the Lebesgue theorem applied to  $\chi_A$  gives that for all  $x_0 \in \text{int}_\mu(A) \setminus S_U^\mu$  and all  $\varepsilon > 0$  small, there is a  $\delta > 0$ : for all  $r < \delta$ ,  $\mu(\mathbb{B}_r(x_0) \cap A) \geq (1 - \varepsilon) \mu(\mathbb{B}_r(x_0))$ . Consequently, since  $U^\mu = U$   $\mu$ -a.e.,

$$\begin{aligned} \mu(A \cap \{|U - u_0| < 2\varepsilon\}) &\geq \mu(\{|U^\mu(x_0) - u_0| < \varepsilon\} \cap \{|U - U^\mu(x_0)| < \varepsilon\} \cap A) \\ &\geq \mu(\mathbb{B}_r(x_0) \cap \{|U - U^\mu(x_0)| < \varepsilon\} \cap A) \\ &\geq (1 - \varepsilon) \mu(\mathbb{B}_r(x_0) \cap A) \\ &\geq (1 - \varepsilon)^2 \mu(\mathbb{B}_r(x_0)) > 0. \end{aligned}$$

Thus, we obtain that  $\mu \llcorner A - \text{ess inf } |U - u_0| = 0$  as desired.  $\square$

**Remark 5.7.** The weak\* limits of  $\mathcal{F}_{1+\frac{1}{m}}^{C,m}$  and  $\mathcal{F}^{C,m}$  in the proof of Lemma 4.5 can actually be taken simultaneously. In view of this, we collect all the ingredients that we will utilize in the sequel:

$$\left\{ \begin{array}{l} \text{If } C \subseteq \mathbb{R}^N \text{ is a compact convex set with } 0 \in \text{int}(C), \text{ we take} \\ \\ C_m := \left\{ \text{dist}(\cdot, C) \leq \frac{1}{m} \right\} \subseteq \mathbb{R}^N, \quad m = 1, 2, \dots \\ \\ \text{Each } C_m \text{ is smooth verifying (O1) and } C_m \searrow C. \text{ By Lemma 4.4, there} \\ \text{exists a non-expansive locally Lipschitz Folding Map satisfying} \\ \\ \mathcal{F}_m^C := \mathcal{F}_{1+\frac{1}{m}}^{C,m} : \mathbb{R}^N \longrightarrow \mathbb{R}^N, \quad m = 1, 2, \dots, \\ (\mathcal{F}_m^C)_1^\infty \subseteq [W_{loc}^{1,\infty}(\mathbb{R}^N)]^N, \quad \text{Lip}[\mathcal{F}_m^C] = 1, \\ \\ \begin{cases} \mathcal{F}_m^C(C_m \cap E) = C_m \cap E, & \text{for any } E \subseteq \mathbb{R}^N, \\ \mathcal{F}_m^C(B_m \setminus C_m) \subseteq \left(1 + \frac{1}{m}\right) C_m \setminus \text{int}(C_m). \end{cases} \\ \\ \text{By Lemma 4.4, there exist reflections } R_1, \dots, R_{r(m)} \text{ in } \mathcal{R}(C), u_1, \dots, u_{r(m)} \\ \text{in } \mathbb{R}^N \text{ such that } \mathcal{F}_m^C \text{ is composed by Affine reflections in } \text{AO}(N, \mathbb{R}): \\ \\ \mathcal{F}_m^C(u) = \left( ((uR_1 + u_1) \cdots) R_{r(m-1)} + u_{r(m-1)} \right) R_{r(m)} + u_{r(m)}. \\ \\ \text{For } m \geq k, k \text{ fixed, we have } \|\mathcal{F}_m^C\|_{[W_{loc}^{1,\infty}(B_k)]^N} \leq 2 \max \{|u| : u \in C_1\} + 1. \\ \text{Hence, utilizing Lemma 4.5, if } \mathcal{P}^C : \mathbb{R}^N \longrightarrow C \text{ is the projection on } C, \\ \\ \mathcal{F}_m^C \xrightarrow{*} \mathcal{P}^C \text{ in } [W_{loc}^{1,\infty}(\mathbb{R}^N)]^N, \text{ as } m \rightarrow \infty, \\ \\ \text{up to a subsequence. If } C \text{ is an obtuse simplex satisfying (O), then} \\ (R_j)_1^{r(m)} \text{ can be chosen in the finite reduced set } \mathcal{R}^*(C). \end{array} \right. \quad (\diamond)$$

We record here a few technical facts which are needed in the proof of Lemma 4.6.

**Properties of (Essrng):** If  $U \in [L^\infty(A, \mu)]^N$  with  $\mu$  Borel satisfying  $\mu(E) > 0$  for all open  $E \neq \emptyset$ , then  $\mu$ -ess  $U(A)$  is compact and it can be proved that at least one point on  $\mu$ -ess  $U(A)$  realizes  $\|U\|_{[L^\infty(A, \mu)]^N}$ :

$$\|U\|_{[L^\infty(A, \mu)]^N} = \max_{u \in \mu\text{-ess } U(A)} |u|. \quad (\text{essrng})$$

**Properties of functions in (AS):** Since  $[W_0^{1;p,q}(\Omega)]^N \subseteq [W_{loc}^{1,\min\{p,q\}}(\mathbb{R}^n)]^N$ , if  $n < \min\{p, q\}$ ,  $[W_0^{1;p,q}(\Omega)]^N$  consists of equivalence classes of  $\mathcal{L}^n \llcorner \Omega$ -a.e. functions whose representatives are in  $C^{1+(1-\frac{n}{\min\{p,q\}})}(\Omega)$ . For  $n \geq \min\{p, q\}$ , since  $[W_0^{1;p,q}(\Omega)]^N \subseteq [W_{loc}^{1,1}(\mathbb{R}^n)]^N \subseteq [BV_{loc}(\mathbb{R}^n)]^N$ , the representatives in  $[W_{loc}^{1;p,q}(\mathbb{R}^n)]^N$  are defined on  $\mathbb{R}^n$  except for an  $\mathcal{H}^{n-1}$ -nullset (Evans-Gariepy [E-G]).

Functions in  $[W^{1,\infty}(\Omega)]^N$  will always be tacitly identified with their representatives.

**Proof of Lemma 4.6.** Since  $C$  is a fixed convex set, we drop superscripts. Proof of (C1): let first  $p = \infty$ . By ( $\diamond$ ), we have  $\mathcal{F}_m \rightarrow \mathcal{P}$  in  $[L_{loc}^\infty(\mathbb{R}^N)]^N$ . Since  $U \in [L^\infty(\Omega)]^N$ , (T) implies that  $U = U^m$  for all  $m \geq \|U\|_{[L^\infty(\Omega)]^N}$ . By Remark 4.9, we have

$$\begin{aligned} \|\mathcal{F}_m(U) - \mathcal{P}(U)\|_{[L^\infty(\Omega)]^N} &= \text{ess sup}_{\Omega \setminus S_U^*} |\mathcal{F}_m(U^*) - \mathcal{P}(U^*)| \\ &= \sup_{\Omega \setminus S_U^*} |\mathcal{F}_m(U^*) - \mathcal{P}(U^*)|. \end{aligned}$$

Hence, utilizing the representation formulas (EssRng) and (essrng), we have

$$\begin{aligned} \|\mathcal{F}_m(U) - \mathcal{P}(U)\|_{[L^\infty(\Omega)]^N} &= \sup_{u \in U^*(\Omega \setminus S_U^*)} |\mathcal{F}_m(u) - \mathcal{P}(u)| \\ &\leq \max_{u \in \text{ess } U(\Omega)} |\mathcal{F}_m(u) - \mathcal{P}(u)| \\ &\leq \max_{|u| \leq \text{ess sup}_\Omega |U|} |\mathcal{F}_m(u) - \mathcal{P}(u)| \\ &= \|\mathcal{F}_m - \mathcal{P}\|_{[L^\infty(\{u \in \mathbb{R}^N : |u| \leq \|U\|_{[L^\infty(\Omega)]^N}\})]^N}. \end{aligned}$$

Thus (C1) has been established when  $p = \infty$ . Let now  $p \in [1, \infty)$ . In view of (T), we have the bound  $|U^m - U| \leq 2|U| \chi_{\{|U| \geq m\}}$ , thus  $U^m \rightarrow U$   $\mathcal{L}^n \llcorner \Omega$ -a.e. and in  $[L^p(\Omega)]^N$ . By ( $\diamond$ ) and (T) we have for large  $m$  and  $\mathcal{L}^n \llcorner \Omega$ -a.e., that

$$\begin{aligned} |\mathcal{F}_m(U^m) - \mathcal{P}(U)|^p &\leq 2^{p-1} (|\mathcal{F}_m(U^m)|^p + |\mathcal{P}(U)|^p) \\ &\leq 2^{p-1} (|\mathcal{F}_m(U^m)|^p \chi_{\{U \in C_m\}} + |\mathcal{F}_m(U^m)|^p \chi_{\{U \notin C_m\}} + |U|^p) \\ &\leq 2^{p-1} (|U|^p \chi_{\{U \in C_1\}} + 2^p (\text{diam}(C))^p \chi_{\{U \notin C\}} + |U|^p), \end{aligned}$$

which gives the pointwise estimates

$$\begin{cases} |\mathcal{F}_m(U^m)|^p \leq |U|^p + 2^p (\text{diam}(C))^p \chi_{\{U \notin C\}}, \\ |\mathcal{F}_m(U^m) - \mathcal{P}(U)|^p \leq 2^p |U|^p + 2^{2p-1} (\text{diam}(C))^p \chi_{\{U \notin C\}}. \end{cases} \quad (\text{B1})$$

Recall that  $|U| \in L^p(\Omega)$  and by assumption of Lemma 4.6  $\mathcal{L}^n(\{U \notin C\}) < \infty$ . Now we utilize (B1) and that  $\mathcal{F}_m(U^m) \rightarrow \mathcal{P}(U)$   $\mathcal{L}^n \llcorner \Omega$ -a.e.. The latter follows as a consequence of ( $\diamond$ ) and the estimate

$$|\mathcal{F}_m(U^m) - \mathcal{P}(U)| \leq 4 \text{diam}(C) \chi_{\{|U| \geq m\}} + |\mathcal{F}_m(U) - \mathcal{P}(U)|. \quad (\text{B2})$$



Utilizing the Dominated Convergence theorem, we see that (C1) is established when  $p \in [1, \infty)$ .

Proof of (C2a), (C2b), (C2c): Upon differentiating (T) weakly, we obtain  $|\nabla U^m| \leq 2\sqrt{N}|\nabla U|$   $\mathcal{L}^n \llcorner \Omega$ -a.e.. By  $(\diamond)$ , there are  $R_1, \dots, R_{r(m)} \in O(N, \mathbb{R})$  such that  $|\nabla(\mathcal{F}_m(U^m))| = |\nabla U^m R_1 \cdots R_{r(m)}| = |\nabla U^m|$ . Thus, we obtain:

$$|\nabla(\mathcal{F}_m(U^m))| = |\nabla U^m| \leq 2\sqrt{N}|\nabla U|, \quad (\text{B3})$$

at least  $\mathcal{L}^n \llcorner \Omega$ -a.e.. Assume first that  $\min\{p, q\} > 1$ . Then, by definition of (AfS), the weak topology  $\mathcal{T}_{1;p,q}^w$  on  $[W_0^{1;p,q}(\Omega)]^N$  has (i) weak-weak, (ii) weak-weak\*, (iii) weak\*-weak\* compactness depending on whether (i)  $\max\{p, q\} < \infty$ , (ii)  $\min\{p, q\} < \infty = \max\{p, q\}$ , (iii)  $p = q = \infty$ . The bounds (B1), (B3) imply the estimates

$$\|\mathcal{F}_m(U^m)\|_{[L^p(\Omega)]^N} \leq \|U\|_{[L^p(\Omega)]^N} + 2 \operatorname{diam}(C) \mathcal{L}^n(\{U \notin C\}), \quad (\text{B4(i)})$$

$$\|\nabla(\mathcal{F}_m(U^m))\|_{[L^q(\Omega)]^{n \times N}} \leq 2\sqrt{N} \|\nabla U\|_{[L^q(\Omega)]^{n \times N}}. \quad (\text{B4(ii)})$$

By  $(\diamond)$ , since  $0 \in \operatorname{int}(C)$ ,  $\mathcal{F}_m$  does not affect the zero trace values of  $U$ , leaving the subspace  $[W_0^{1;p,q}(\Omega)]^N$  invariant:  $\mathcal{F}_m([W_0^{1;p,q}(\Omega)]^N) \subseteq [W_0^{1;p,q}(\Omega)]^N$ . Thus, there is a  $V \in [W_0^{1;p,q}(\Omega)]^N$  such that up to a certain subsequence

$$\mathcal{F}_m(U^m) \rightharpoonup V \text{ weakly with respect to } \mathcal{T}_{1;p,q}^w, \text{ in } [W_0^{1;p,q}(\Omega)]^N.$$

Since  $[W_0^{1;p,q}(\Omega)]^N \subseteq [W_{\operatorname{loc}}^{1,\min\{p,q\}}(\mathbb{R}^n)]^N$ , by standard local strong compactness reasoning

$$\mathcal{F}_m(U^m) \rightarrow V \text{ in } [L_{\operatorname{loc}}^{\min\{p,q\}}(\mathbb{R}^n)]^N, \text{ and } \mathcal{L}^n \llcorner \Omega - \text{a.e..}$$

Since  $\mathcal{F}_m(U^m) \rightarrow \mathcal{P}(U)$  in  $[L^p(\mathbb{R}^n)]^N \subseteq [L_{\operatorname{loc}}^{\min\{p,q\}}(\mathbb{R}^n)]^N$ , we obtain  $V = \mathcal{P}(U)$   $\mathcal{L}^n \llcorner \Omega$ -a.e..

Assume now that  $\min\{p, q\} = 1$ . Then,  $[W_0^{1;p,q}(\Omega)]^N \subseteq [W_{\operatorname{loc}}^{1,1}(\Omega)]^N$ . We define the Radon measures

$$\begin{cases} \mu_m := \mathcal{L}^n \llcorner \mathcal{F}_m(U^m) & \in [\mathcal{M}(\Omega)]^N, \\ \nu_m := \mathcal{L}^n \llcorner \nabla(\mathcal{F}_m(U^m)) & \in [\mathcal{M}(\Omega)]^{n \times N}, \quad m = 1, 2, \dots \end{cases} \quad (\text{B5})$$

By (B1) for  $p = 1$  and (B3) we have the following total variation estimates for the measures (B5)

$$\begin{cases} \|\mu_m\| \leq \mathcal{L}^n \llcorner |U| + 2 \operatorname{diam}(C) \mathcal{L}^n \llcorner \{U \notin C\}, \\ \|\nu_m\| \leq 2\sqrt{N} \mathcal{L}^n \llcorner |\nabla U|. \end{cases} \quad (\text{B6})$$

By local weak\* compactness of the space of the Radon measures and (B6), there exist measures  $\mu \in [\mathcal{M}(\Omega)]^N$  and  $\nu \in [\mathcal{M}(\Omega)]^{n \times N}$  such that  $\mu_m \xrightarrow{*} \mu$  and  $\nu_m \xrightarrow{*} \nu$  as  $m \rightarrow \infty$ . By weak\* LSC of the total variation, upon passing to the weak\* limit in (B6), we have for any  $A \subseteq \Omega$  open that

$$\|\mu\|(A) \leq \liminf_{m \rightarrow \infty} \|\mu_m\|(A) \leq \int_A |U| d\mathcal{L}^n + 2 \operatorname{diam}(C) \mathcal{L}^n(A \cap \{U \notin C\}), \quad (\text{B7(i)})$$

$$\|\nu\|(A) \leq \liminf_{m \rightarrow \infty} \|\nu_m\|(A) \leq 2\sqrt{N} \int_A |\nabla U| d\mathcal{L}^n. \quad (\text{B7(ii)})$$

By Radon regularity, (B7(i)) and (B7(ii)) imply that  $\|\mu\| \ll \mathcal{L}^n$  and  $\|\nu\| \ll \mathcal{L}^n$ . Thus, there exist  $V \in [L_{\operatorname{loc}}^1(\Omega)]^N$  and  $W \in [L_{\operatorname{loc}}^1(\Omega)]^{n \times N}$  such that

$$\begin{cases} \mathcal{F}_m(U^m) \rightarrow V & \text{in } [L_{\operatorname{loc}}^1(\Omega)]^N, \\ \nabla(\mathcal{F}_m(U^m)) \rightarrow W & \text{in } [L_{\operatorname{loc}}^1(\Omega)]^{n \times N}. \end{cases} \quad (\text{B8})$$

By weak closure of the graph of  $\nabla : [W^{1,1}(\Omega')]^N \subseteq [L^1(\Omega')]^N \rightarrow [L^1(\Omega')]^{n \times N}$  for all  $\Omega' \subset \subset \Omega$  and local strong compactness, we get  $W = \nabla V$  and  $\mathcal{F}_m(U^m) \rightarrow V$  in  $[L^1_{\text{loc}}(\Omega)]^N$ . Since  $\mathcal{F}_m(U^m) \rightarrow \mathcal{P}(U)$  in  $[L^p(\Omega)]^N$ , we get  $V = \mathcal{P}(U)$ . By weak/weak\* LSC of the  $L^p$  and  $L^q$  norms ( $1 \leq p, q \leq \infty$ ), passing to the limit to (B4(i)), (B4(ii)) and utilizing that  $[W^{1,p,q}_0(\Omega)]^N$  is  $\mathcal{T}^w_{1;p,q}$ -weakly closed, we obtain that  $\mathcal{P}(U) \in [W^{1,p,q}_0(\Omega)]^N$  and  $\mathcal{F}_m(U^m) \rightarrow \mathcal{P}(U)$  weakly with respect to  $\mathcal{T}^w_{1;p,q}$ . Lemma (4.6) follows.  $\square$

We present first the proof of Theorem 3.7 which is easier and next that of Theorem 3.1.

**Proof of Theorem 3.7.** Let  $U = U^*$  be an absolute minimizer in  $[W^{1,\infty}_{U_0}(\Omega)]^N$  of  $(E_\infty)$  verifying (AM). Up to a translation and a dimensional reduction  $\mathbb{R}^N \rightsquigarrow \mathbb{R}^{\dim(\text{int}(C))}$ , we may assume that  $0 \in \text{int}(C) \neq \emptyset$ . Let  $\mathcal{F}_m : \mathbb{R}^N \rightarrow \mathbb{R}^N$  be the  $m$ -Folding Map of  $(\diamond)$  and  $\mathcal{P} : \mathbb{R}^N \rightarrow C$  the projection, both referring to the fixed convex set  $C$ . By (C1) of Lemma 4.6, we have  $\mathcal{F}_m(U) \rightarrow \mathcal{P}(U)$  in  $[L^\infty(\Omega)]^N$  as  $m \rightarrow \infty$ . By the assumption (HR) and  $(\diamond)$ , we have for any  $\Omega' \subseteq \Omega$ ,  $m \geq \|U\|_{[L^\infty(\Omega)]^N}$

$$\begin{aligned} E_\infty(\mathcal{F}_m(U), \Omega') &= \text{ess sup}_{\Omega'} \mathcal{L}(\nabla(\mathcal{F}_m(U)), \mathcal{F}_m(U), -) \\ &= \text{ess sup}_{\Omega'} \mathcal{L}(\nabla U R_1 \cdots R_{r(m)}, \mathcal{F}_m(U), -) \\ &= \text{ess sup}_{\Omega'} \mathcal{L}(\nabla U, \mathcal{F}_m(U), -). \end{aligned}$$

Again by  $(\diamond)$  and (C1), for any  $\Omega' \subseteq \Omega$  we have  $\mathcal{F}_m(U(\Omega')) \subseteq 4C$ . Thus, upon replacing  $\nabla U$  by its  $\mathcal{L}^n$ -precise representative and since  $\Omega' = \text{int}_{\mathcal{L}^n}(\Omega')$  (see (EssRng)), assumption (HL) implies

$$\begin{aligned} &\left| E_\infty(\mathcal{F}_m(U), \Omega') - \text{ess sup}_{\Omega'} \mathcal{L}(\nabla U, \mathcal{P}(U), -) \right| \\ &\leq \text{ess sup}_{\Omega'} \left| \mathcal{L}(\nabla U, \mathcal{F}_m(U), -) - \mathcal{L}(\nabla U, \mathcal{P}(U), -) \right| \\ &\leq \left\{ \sup_{|P| \leq \|\nabla U\|_{[L^\infty(\Omega)]^{n \times N}}} \left( \text{ess sup}_{x \in \Omega'} ([\mathcal{L}(P, -, x)]_{C^{0+\gamma}(4C)}) \right) \right\} \|\mathcal{F}_m(U) - \mathcal{P}(U)\|_{[L^\infty(\Omega)]^N}^\gamma. \end{aligned}$$

As a result of (C1), for any  $\Omega' \subseteq \Omega$  we obtain


$$\lim_{m \rightarrow \infty} E_\infty(\mathcal{F}_m(U), \Omega') = \text{ess sup}_{\Omega'} \mathcal{L}(\nabla U, \mathcal{P}(U), -). \quad (\text{Lim})$$

Let  $\varepsilon > 0$  be arbitrary but fixed. We will show that for any  $\varepsilon > 0$ ,  $U(\Omega) \subseteq (1 + \varepsilon)C$ . To this end, we will utilize the assumption (HC),  $(\diamond)$  and Lemma 4.6 to construct an  $(E_\infty)$ -energy comparison function. We assume that  $(FP_\infty \text{ (I)})$  fails, otherwise we are done. Hence,

$$U(\Omega) \not\subseteq C.$$

Since  $U \in [W^{1,\infty}_{U_0}(\Omega)]^N$ , there is an  $\varepsilon_\infty \in (0, \infty)$  such that (recall (EssCo), (Co) and (CO)):

$$\begin{cases} U(\Omega) \subseteq (1 + \varepsilon_\infty)C, \\ \Omega_t := \{x \in \Omega : U(x) \notin (1 + t)C\} \neq \emptyset, & \text{for all } 0 < t < \varepsilon_\infty, \\ U(\Omega) \cap (1 + t)C = \emptyset, & \text{for all } t > \varepsilon_\infty. \end{cases}$$



By continuity and **(CO)**,  $\Omega_t \subset \subset \Omega$ . We may assume  $\varepsilon < \varepsilon_\infty$  and we choose  $t := \varepsilon$ . By **(Lim)**, for every  $\sigma > 0$ , there is an  $m = m(\sigma) \in \mathbb{N}$  such that for all  $m > m(\sigma)$ ,

$$E_\infty(\mathcal{F}_m(U), \Omega_\varepsilon) \leq \operatorname{ess\,sup}_{\Omega_\varepsilon} \mathcal{L}(\nabla U, \mathcal{P}(U), \cdot) + \sigma. \quad (\text{F1})$$

Since  $U(\Omega_\varepsilon) \subseteq \mathbb{R}^N \setminus (1 + \varepsilon)C \subseteq \mathbb{R}^N \setminus C$  and  $\mathcal{P}(U(\Omega_\varepsilon)) \subseteq \partial C$ , for every  $x \in \Omega_\varepsilon \setminus S_{\nabla U}^*$ , we have as a result of **(HC)** that

$$\mathcal{L}(\nabla U(x), \mathcal{P}(U(x)), x) < \mathcal{L}(\nabla U(x), U(x), x)$$

and since  $\varepsilon > 0$ , that  $\operatorname{dist}(\overline{U(\Omega_\varepsilon)}, \partial C) > 0$ . Consequently,

$$\sigma(\varepsilon) := E_\infty(U, \Omega_\varepsilon) - \operatorname{ess\,sup}_{\Omega_\varepsilon} \mathcal{L}(\nabla U, \mathcal{P}(U), \cdot) > 0. \quad (\text{F2})$$

In fact,  $\sigma(t) > 0$  for all  $t \in (0, \varepsilon_\infty)$ . For  $\sigma := \frac{1}{2}\sigma(\varepsilon)$ , there is an  $m(\varepsilon) \in \mathbb{N}$  which may be chosen to satisfy

$$m(\varepsilon) > \max \left\{ \frac{1 + c_{\min}}{2}, \frac{6}{c_{\min}\varepsilon} \right\}, \quad c_{\min} := \min \{|u| : u \in \partial C\} \quad (\text{m})$$

such that for all  $m \geq m(\varepsilon)$ , by equations **(F1)** and **(F2)**,

$$\begin{aligned} E_\infty(\mathcal{F}_m(U), \Omega_\varepsilon) &\leq \frac{\sigma(\varepsilon)}{2} + \operatorname{ess\,sup}_{\Omega_\varepsilon} \mathcal{L}(\nabla U, \mathcal{P}(U), \cdot) \\ &< E_\infty(U, \Omega_\varepsilon). \end{aligned} \quad (\spadesuit)$$

By **(m)**, the choice of  $m(\varepsilon)$  together with **(\diamond)** implies that for all  $m \geq m(\varepsilon)$ ,

$$\begin{aligned} \mathcal{F}_m(U(\Omega)) &\subseteq \left(1 + \frac{1}{m}\right) \left\{ \operatorname{dist}(\cdot, C) \leq \frac{1}{m} \right\} \\ &\subseteq \left(1 + \frac{1}{m}\right) \left(1 + \frac{1}{c_{\min}m}\right) C \\ &= \left[1 + \frac{1}{m} \left(\frac{1}{m} + \frac{1}{c_{\min}} + \frac{1}{c_{\min}m}\right)\right] C \\ &\subseteq \left(1 + \frac{3}{m c_{\min}}\right) C \\ &\subseteq \left(1 + \frac{\varepsilon}{2}\right) C. \end{aligned} \quad (\text{F3})$$

Thus, we obtain

$$\mathcal{F}_m(U(\Omega)) \subseteq (1 + \varepsilon) C. \quad (\spadesuit\spadesuit)$$

By assumption,  $\overline{\text{co}}(U_0(\partial\Omega)) \subseteq C$ . Upon recalling (Co) and (CO), for any  $\delta > 0$ , there is an  $A_\delta \subseteq \overline{\Omega}$  relatively open such that

$$\overline{\text{co}}(U_0(A_\delta)) \subseteq \left\{ \text{dist}(\cdot, \overline{\text{co}}(U_0(\partial\Omega))) \leq \delta \right\}.$$

Again by  $(\diamond)$ ,

$$\mathcal{F}_m \left( \left\{ \text{dist}(\cdot, \overline{\text{co}}(U_0(\partial\Omega))) \leq \frac{1}{m} \right\} \right) \subseteq \left\{ \text{dist}(\cdot, \overline{\text{co}}(U_0(\partial\Omega))) \leq \frac{1}{m} \right\}.$$

For  $\delta(m) := \frac{1}{2m}$ , we see that

$$U_0(A_{\frac{1}{2m}}) \subseteq \overline{\text{co}}(U_0(A_{\frac{1}{2m}})) \subseteq \left\{ \text{dist}(\cdot, C) \leq \frac{1}{2m} \right\}.$$

This implies that the nonlinear operator  $\mathcal{F}_m$  leaves the affine subspace  $[W_{U_0}^{1,\infty}(\Omega)]^N$  invariant, since  $A_{\frac{1}{2m}} \subseteq \overline{\Omega} \setminus \left\{ \text{dist}(U_0, C) \leq \frac{1}{2m} \right\}$ . We obtain

$$\mathcal{F}_m(U(\Omega)) \in [W_{U_0}^{1,\infty}(\Omega)]^N \quad \text{and} \quad \overline{\text{co}}(\mathcal{F}_m(U(\partial\Omega))) = \overline{\text{co}}(U(\partial\Omega)). \quad (\spadesuit\spadesuit\spadesuit)$$

Putting  $(\spadesuit)$ ,  $(\spadesuit\spadesuit)$ ,  $(\spadesuit\spadesuit\spadesuit)$  together we have that all deformations  $\mathcal{F}_m(U)$  for  $m > m(\varepsilon)$  given by (m) comprise comparison functions with  $E_\infty(\mathcal{F}_m(U), \Omega_\varepsilon) < E_\infty(U, \Omega_\varepsilon)$ ,  $\Omega_\varepsilon \subset\subset \Omega$ . In view of (AM), the absolute minimizer  $U$  necessarily satisfies

$$U(\Omega) \subseteq (1 + \varepsilon) C,$$

for all  $\varepsilon > 0$ . The desired localization conclusion ( $FP_\infty$  (I)) of theorem 3.7 follows.  $\square$

**Proof of Theorem 3.1.** Let  $U = U_i$  be the term of a minimizing sequence of  $(E_1)$  in  $[W_{U_0}^{1,p,q}(\Omega)]^N$ , with the exponent  $q$  as in the assumption (HB). By assumption, (recall (EssCo), (Co), (CO)) we have  $\text{app } \overline{\text{co}}(U_0(\partial\Omega)) \subseteq C$ . We may again assume that  $0 \in \text{int}(C) \neq \emptyset$ . Let  $\mathcal{F}_m, \mathcal{P} : \mathbb{R}^N \rightarrow \mathbb{R}^N$  be the  $m$ -Folding Map of  $(\diamond)$  and the projection, both referring to  $C$  of assumptions (HR), (HC). By (C1) of Lemma 4.6 and the finiteness assumption  $\mathcal{L}^n(\Omega) < \infty$ , we have  $\mathcal{F}_m(U^m) \rightarrow \mathcal{P}(U)$  in  $[L^p(\Omega)]^N$  and  $\mathcal{L}^n \llcorner \Omega$ -a.e. as  $m \rightarrow \infty$ .  $U^m$  is the  $m$ -truncate of  $U$  as in (T). By the assumption (HR) and  $(\diamond)$ , for any  $A \subseteq \Omega$ ,  $\mathcal{L}^n(A) > 0$ , we have

$$\begin{aligned} E_1(\mathcal{F}_m(U^m), A) &= \int_A \mathcal{L}(\nabla(\mathcal{F}_m(U^m)), \mathcal{F}_m(U^m), x) dx \\ &= \int_A \mathcal{L}(\nabla U^m R_1 \cdots R_{r(m)}, \mathcal{F}_m(U^m), x) dx \\ &= \int_A \mathcal{L}(\nabla U^m, \mathcal{F}_m(U^m), x) dx. \end{aligned}$$

We set  $E_m := \{|U| > m\}$ . Upon identifying  $U = U^*$  and  $\nabla U = (\nabla U)^*$ , in view of (T) we have  $\nabla U = \nabla U^m$  everywhere on  $\Omega \setminus E_m \setminus S_{\nabla U}^*$ . In view of (C1),  $(\diamond)$  and Remark 4.9, for large  $m \in \mathbb{N}$  we have  $\mathcal{F}_m(U^m(x))$ ,

$\mathcal{P}(U(x)) \in 4C$ , for all  $x \in \Omega \setminus S_U^*$ . Utilizing the assumptions **(HL)**, **(HL1)**, we estimate:

$$\begin{aligned} \left| E_1(\mathcal{F}_m(U^m), A) - \int_A \mathcal{L}(\nabla U, \mathcal{P}(U), x) dx \right| &\leq \\ &\leq \int_A \left| \mathcal{L}(\nabla U^m, \mathcal{F}_m(U^m), x) - \mathcal{L}(\nabla U, \mathcal{P}(U), x) \right| dx \\ &\leq \int_{A \setminus E_m} \left\{ \sup_{|P| \leq m} \operatorname{ess\,sup}_{x \in A} \left( \operatorname{Lip}[\mathcal{L}(P, \cdot, x), 4C] \right) \right\} |\mathcal{F}_m(U^m) - \mathcal{P}(U)| d\mathcal{L}^n \\ &\quad + \int_{E_m \cap A} \left\{ \mathcal{L}(\nabla U^m, \mathcal{F}_m(U^m), x) + \mathcal{L}(\nabla U, \mathcal{P}(U), x) \right\} dx \end{aligned}$$

(by assumption **(HB)** and **(B3)**)

$$\begin{aligned} &\leq \sup_{|P| \leq m} \operatorname{ess\,sup}_{x \in A} \left( \operatorname{Lip}[\mathcal{L}(P, \cdot, x), 4C] \right) \mathcal{L}^n(A)^{1-\frac{1}{p}} \|\mathcal{F}_m(U^m) - \mathcal{P}(U)\|_{[L^p(A)]^N} \\ &\quad + \int_{E_m} \left\{ C((2\sqrt{N})^q + 1) |\nabla U|^q + 2 \sup_{\xi \in 4C} \beta(\xi, \cdot) \right\} d\mathcal{L}^n. \end{aligned}$$

We conclude:

$$\begin{aligned} \left| E_1(\mathcal{F}_m(U^m), A) - \int_A \mathcal{L}(\nabla U, \mathcal{P}(U), x) dx \right| &\leq \\ &\leq \sup_{P \in \mathbb{R}^{n \times N}} \operatorname{ess\,sup}_{x \in \Omega} \left( \operatorname{Lip}[\mathcal{L}(P, \cdot, x), 4C] \right) \mathcal{L}^n(\Omega)^{1-\frac{1}{p}} \|\mathcal{F}_m(U^m) - \mathcal{P}(U)\|_{[L^p(\Omega)]^N} \\ &\quad + C((2\sqrt{N})^q + 1) \|\nabla U\|_{[L^q(E_m)]^{n \times N}}^q + 2 \left\| \sup_{\xi \in 4C} \beta(\xi, \cdot) \right\|_{L^1(E_m)}. \end{aligned}$$

As a result of **(C1)**, for any  $A \subseteq \Omega$  measurable we obtain

$$\lim_{m \rightarrow \infty} E_1(\mathcal{F}_m(U^m), A) = \int_A \mathcal{L}(\nabla U, \mathcal{P}(U), x) dx \quad (\text{Lim1})$$

$A$ -uniformly. Now we argue by contradiction as in Theorem 3.7. To this end, we assume  $\operatorname{ess} U(\Omega) \not\subseteq C$ , otherwise the conclusion **(FP<sub>1</sub> (I))** follows. Since  $\Omega$  is open,  $\Omega = \operatorname{int}_{\mathcal{L}^n}(\Omega)$  and the formula **(EssRng)** implies

$$\operatorname{ess} U(\Omega) = \overline{U(\Omega \setminus S_U^*)} \not\subseteq C.$$

Necessarily, there exists a measurable set  $A \subseteq \Omega$  with  $\mathcal{L}^n(A) > 0$  such that

$$\operatorname{ess} U(A) = \overline{U(\operatorname{int}_{\mathcal{L}^n}(A) \setminus S_U^*)} \subseteq \mathbb{R}^N \setminus C.$$

We identify as we can  $A \equiv \operatorname{int}_{\mathcal{L}^n}(A)$ , since  $\mathcal{L}^n(A \Delta \operatorname{int}_{\mathcal{L}^n}(A)) = 0$ . Let  $\varepsilon > 0$  be arbitrary but fixed. For all  $x \in A \setminus (S_U^* \cup S_{\nabla U}^*)$ , the relation

$$\mathcal{P}(U(A \setminus S_U^*)) \subseteq \mathcal{P}(\overline{U(A \setminus S_U^*)}) \subseteq \mathcal{P}(\mathbb{R}^N \setminus C) \subseteq \partial C$$

together with **(HC)** imply in view of Remark 4.9 that

$$\mathcal{L}(\nabla U(x), \mathcal{P}(U(x)), x) < \mathcal{L}(\nabla U(x), U(x), x).$$

In addition, since  $\text{dist}(\overline{U(A \setminus S_U^*)}, \partial C) > 0$ , we obtain

$$\sigma(A) := E_1(U, A) - \int_A \mathcal{L}(\nabla U, \mathcal{P}(U), x) dx > 0. \quad (\text{F4})$$

By (Lim1), for  $\sigma := \frac{1}{2}\sigma(A)$ , there is an  $m = m(A) \in \mathbb{N}$  which may be chosen to have as lower bound the right hand side of (m), such that for all  $m \geq m(A)$ , by equation (F4),

$$\begin{aligned} E_1(\mathcal{F}_m(U^m), A) &\leq \frac{\sigma(A)}{2} + \int_A \mathcal{L}(\nabla U, \mathcal{P}(U), x) dx \\ &< E_1(U, A). \end{aligned} \quad (\clubsuit)$$

By our choice of  $m(A)$ , we obtain that (F3), (essrng) and (T) imply for all  $m \geq m(A)$ ,

$$\begin{aligned} \mathcal{F}_m(U^m(\Omega \setminus S_U^*)) &\subseteq \mathcal{F}_m(\overline{U^m(\Omega \setminus S_U^*)}) \\ &= \mathcal{F}_m(\text{ess } U^m(\Omega)) \\ &\subseteq \mathcal{F}_m(\{u \in \mathbb{R}^N : |u| \leq \|U^m\|_{[L^\infty(\Omega)]^N}\}) \\ &= \mathcal{F}_m(\mathbb{B}_m^N(0)) \\ &\subseteq \left(1 + \frac{1}{m}\right) \left\{\text{dist}(\cdot, C) \leq \frac{1}{m}\right\} \\ &\subseteq \left(1 + \frac{\varepsilon}{2}\right) C. \end{aligned} \quad (\clubsuit\clubsuit)$$

By assumption,  $\text{app } \overline{\text{co}}(U_0(\partial\Omega)) \subseteq C$ . Thus, (Co) implies that for  $\delta(m) = \frac{1}{2m}$ , there is an  $A_{\frac{1}{2m}} \subseteq \overline{\Omega}$  relatively open such that

$$\begin{aligned} \text{ess } \overline{\text{co}}(U_0(A_\delta)) &\subseteq \left\{\text{dist}\left(\cdot, \text{app } \overline{\text{co}}(U_0(\partial\Omega))\right) \leq \frac{1}{2m}\right\} \\ &\subseteq \left\{\text{dist}(\cdot, C) \leq \frac{1}{2m}\right\}. \end{aligned}$$

By (◇), the latter set is invariant under  $\mathcal{F}_m$ . By (EssCo) we have

$$\text{ess } U_0\left(A_{\frac{1}{2m}}\right) \subseteq \overline{\text{co}}\left(\text{ess } U_0\left(A_{\delta(m)}\right)\right) = \text{ess } \overline{\text{co}}\left(U_0\left(A_{\frac{1}{2m}}\right)\right).$$

Hence, the nonlinear operator  $\mathcal{F}_m$  maps the affine subspace  $[W_{U_0}^{1;p,q}(\Omega)]^N$  into itself. We obtain

$$\mathcal{F}_m(U^m) \in [W_{U_0}^{1;p,q}(\Omega)]^N \quad \text{and} \quad \text{app } \overline{\text{co}}(\mathcal{F}_m(U^m(\partial\Omega))) = \text{app } \overline{\text{co}}(U(\partial\Omega)). \quad (\clubsuit\clubsuit\clubsuit)$$

Putting (♣), (♣♣), (♣♣♣) together we conclude that all deformations  $\mathcal{F}_m(U^m)$  for  $m > m(A)$  comprise comparison functions whose  $(E_1)$ -minimality implies that  $U$  satisfies

$$\text{ess } U(\Omega) \subseteq (1 + \varepsilon) C$$

for all  $\varepsilon > 0$ . The conclusion (FP<sub>∞</sub> (I)) of theorem 3.1 follows.  $\square$

**Proof of Theorem 3.2.** It is a corollary of Theorem 3.1, utilizing the formula (EssRng) and the  $\mathcal{L}^n$ -a.e. convergence up to a subsequence to the relaxed solution  $U^*$ .  $\square$

## References

- [A-G] S. Alexander, M. Ghomi, *The convex hull property and topology of hypersurfaces with nonnegative curvature*, Advances in Mathematics Volume 180, Issue 1, 2003, 324 - 354.
- [A-F] N. Alikakos, G. Fusco, *On the connection problem for potentials with several global minima*, Ind. J. of Math., J. 57 No. 3 (2008), (to appear).
- [A-F2] N. Alikakos, G. Fusco, *On an Elliptic System with Symmetric Potential and Two Global Minima.*, submitted preprint.
- [A-F-P] L. Ambrosio, N. Fusco, D. Pallara, *Functions of Bounded Variation and Free Discontinuity Problems*, Oxford Mathematical Monographs, Clarendon Press, 2000.
- [Ar1] G. Aronsson, *Minimization problems for the functional  $\sup_x F(x, f(x), f'(x))$* , Arkiv für Mat. 6 (1965), 33 - 53.
- [Ar2] G. Aronsson, *Minimization problems for the functional  $\sup_x F(x, f(x), f'(x))$  II*, Arkiv für Mat. 6 (1966), 409 - 431.
- [Ar3] G. Aronsson, *Extension of functions satisfying Lipschitz conditions*, Arkiv für Mat. 6 (1967), 551 - 561.
- [Ar4] G. Aronsson, *On the partial differential equation  $u^2 x u_{xx} + 2u_x u_y u_{xy} + u^2 y u_{yy} = 0$* , Arkiv für Mat. 7 (1968), 395 - 425.
- [Ar5] G. Aronsson, *Minimization problems for the functional  $\sup_x F(x, f(x), f'(x))$  III*, Arkiv für Mat. (1969), 509 - 512.
- [Ar6] G. Aronsson, *On certain singular solutions of the partial differential equation  $u^2 x u_{xx} + 2u_x u_y u_{xy} + u^2 y u_{yy} = 0$* , Manuscripta Math. 47 (1984), no 1-3, 133 - 151.
- [A-C-J] G. Aronsson, M. Crandall, P. Juutinen *A tour of the theory of absolutely minimizing functions*, Bulletin of the AMS, New Series, 41 (2004), 439 - 505.
- [Ba] E. N. Barron, *Viscosity solutions and analysis in  $L^\infty$* , in *Nonlinear Analysis, Differential Equations and Control*, Dordrecht 1999, 160.
- [B-E-J] E. N. Barron, L. C. Evans, R. Jensen, *The Infinity Laplacian, Aronssons Equation and their Generalizations*, Transactions of the AMS, Vol. 360, Nr 1, Jan 2008, electr/ly published on July 25, 2007.
- [BJW1] E. N. Barron, R. Jensen and C. Wang, *The Euler equation and absolute minimizers of  $L^\infty$  functionals*, Arch. Rational Mech. Analysis 157 (2001), 255 - 283.
- [BJW2] E. N. Barron, R. Jensen and C. Wang, *Lower Semicontinuity of  $L^\infty$  Functionals* Ann. I. H. Poincaré AN 18, 4 (2001) 495 - 517.
- [B-F1] M. Bildhauer, M. Fuchs, *Partial regularity for a class of anisotropic variational integrals with convex hull property*, J. Asymptotic Analysis, Vol. 32, Nr 3-4, 293 - 315, 2002.
- [B-F2] M. Bildhauer, M. Fuchs, *A geometric maximum principle for variational problems in spaces of vector valued functions of bounded variation*, Fachrichtung 6.1 - Preprint No. 211, Univ. des Saarlandes, 2008.
- [Ca] S. Campanato, *A Maximum Principle for Nonlinear Elliptic Systems: Boundary Fundamental Estimates*, Adv. in Math, 66, (1987) 291 - 317.
- [Ce] A. Cellina *On the Strong Maximum Principle*, Proceedings of the AMS, Vol. 130, Nr 2, Pages 413 - 418, 2001.
- [C-M1] T. H. Colding, W. P. Minicozzi II, *An excursion into geometric analysis*, preprint arXiv:math/0309021v1.
- [C-M2] T. H. Colding, W. P. Minicozzi II, *Minimal Submanifolds*, preprint arXiv:math/0504158v2.
- [Cr1] M. Crandall *A Visit with the  $\infty$ -Laplacian*, in *Calculus of Variations and Non-Linear Partial Differential Equations*, Springer Lecture notes in Mathematics 1927, CIME, Cetraro Italy 2005.
- [Cr2] M. Crandall, *An Efficient Derivation of the Aronsson Equation*, Arch. Rational Mech. Anal. 167 (2003) 4, 271 - 279.
- [C-I-L] M. Crandall, H. Ishii, P.-L. Lions, *User's Guide to Viscosity Solutions of 2nd Order Partial Differential Equations*, Bulletin of the AMS, Vol. 27, Nr 1, Pages 1 - 67, 1992.
- [C-W-Y] M. Crandall, C. Wang, Y. Yu, *Derivation of the Aronsson Equation for  $C^1$  Hamiltonians*, Transactions of the AMS, to appear.

- [D] B. Dacorogna, *Direct Methods in the Calculus of Variations*, Applied Mathematical Sciences 78, Springer - Verlag, 2008 edition.
- [D-W] D. Li and L. Wang, *A new maximum principle of elliptic differential equations in divergence form*. Proc. Amer. Math. Soc. 136 2823 - 2828 (2008).
- [D-L-M] D'Ottavio, A., Leonetti, F., Musciano, C., *Maximum principle for vector valued mappings minimizing variational integrals*, Atti Sem. Mat. Fis. Uni. Modena XLVI (1998), 677 - 683.
- [E-G] L.C. Evans, R.F. Gariepy, *Measure Theory and Fine Properties of Functions*, CRC Press, Studies in Advanced Mathematics, 1992.
- [F-M-Q] P. Felmer, M. Montenegro, A. Quaas, *A note on the strong maximum principle and the compact support principle*, JDE (2008), doi:10.1016/j.jde.2008.02.034.
- [F-Q] P. Felmer, A. Quaas, *On the Strong Maximum Principle for Quasilinear Elliptic Equations and Systems*, Advances in Differential Equations, Vol. 7, Nr 1, Jan. 2002, pp 25 - 46.
- [Fr] L. E. Fraenkel, *An Introduction to Maximum Principles and Symmetry in Elliptic Problems*, Cambridge Tracts in Mathematics 128, 2000.
- [F] M. Fuchs, *Existence via partial regularity for degenerate systems of variational inequalities with natural growth*, Comment. Math. Univ. Carolinae 33, 3 (1992), 427 - 435.
- [G-W-Y] R. Gariepy, Ch. Wang, Y. Yu, *Generalized Cone Comparison Principle for Viscosity Solutions of the Aronsson Equation and Absolute Minimizers*, Communications in PDE, 31, 1027 - 1046, 2006.
- [G-T] D. Gilbarg, N. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer, 1998, revised 3rd edition
- [H] E. Hopf, *Elementäre Bemerkungen über die Lösungen Partieller Differentialgleichungen Zweiter Ordnung vom Elliptischen Typus*, Sitzungsberichte Preussische Akademie der Wissenschaften, Berlin, 147 - 152, 1927.
- [Je1] R. Jensen, *The Maximum Principle for Viscosity Solutions of Fully Nonlinear Second Order Partial Differential Equations*, Arch. Rat. Mech. Anal. Vol. 101, Nr 1, 1988.
- [Je2] R. Jensen, *Uniqueness of Lipschitz extensions minimizing the sup-norm of the gradient*, Arch. Rational Mech. Analysis 123 (1993), 51-74.
- [JWY] R. Jensen, C. Wang, Y. Yu, *Uniqueness and Nonuniqueness of Viscosity Solutions to Aronsson's Equation*, Arch. Rational Mech. Analysis, (DOI) 10.2007/10.1007/s00205-007-0093-1.
- [K-A] N. Katzourakis, N. Alikakos, *Heteroclinic Travelling Waves of Gradient Diffusion Systems*, Transactions of the AMS, accepted, <http://arxiv.org/abs/0712.2800>.
- [K-A2] N. Katzourakis, N. Alikakos, *Semilinear Elliptic Systems Describing Diffusive Travelling Waves*, in preparation.
- [Ka] N. Katzourakis, *On the Loss of Compactness of Quasilinear Systems in  $W^1$  & BV Spaces*, in preparation.
- [Le] S. Leonardi, *A Maximum Principle for Linear Elliptic Systems with Discontinuous Coefficients*, Comment. Math. Univ. Carolinae 45 (2004) 457 - 474.
- [L-S] F. Leonetti, F. Siepe, *Maximum Principle for Vector Valued Minimizers*, Journal of Convex Analysis 12 (2005) 267 - 278.
- [M] P. Mattila, *Geometry of Sets and Measures in Euclidean Space, Fractals and Rectifiability*, Cambridge Studies in Advanced Mathematics 44, 2004.
- [N-S] F. Nečas, J. Stará, *Principio di Massimo per Sistemi Ellittici Quasi Lineari non Diagonali*, Bol. Un. Mat. It. 4 (1972) 1-10.
- [Os1] R. Osserman, *The Convex Hull Property of Immersed Manifolds*, J. Diff. Geom. 6 (1971), 267 - 270.
- [Os2] R. Osserman, *A Survey of Minimal Surfaces*, 2nd edition, Dover, New York, 1986.
- [P-W] M. Protter, H. Weinberger, *Maximum Principles in Differential equations*, Englewood Cliffs, N.J., Prentice - Hall, 1967.
- [P-S] P. Puzzi, J. Serrin, *The Maximum Principle*, Progress in Nonlinear Differential Equations and Applications 73, Birkhäuser, 2007.
- [R-S] P. Roselli, B. Sciunzi, *A strong comparison principle for the  $p$ -Laplacian*. Proc. Amer. Math. Soc. 135, 3217-3224 (2008).
- [Y] Y. Yu,  *$L^\infty$  Variational Problems and Aronsson Equations*, Arch. Rational Mech. Anal. 182, 153 - 180, (2006).